

DIFFERENTIAL GEOMETRIC ASPECTS OF CAUSAL STRUCTURES

OMID MAKHMALI

ABSTRACT. This article is concerned with causal structures, which are defined as a field of tangentially non-degenerate projective hypersurfaces in the projectivized tangent bundle of a manifold. The local equivalence problem of causal structures on manifolds of dimension at least four is solved using Cartan's method of equivalence, leading to an $\{e\}$ -structure over some principal bundle. It is shown that these structures correspond to parabolic geometries of type $(D_n, P_{1,2})$ and $(B_n, P_{1,2})$ when $n \geq 4$ and $(D_3, P_{1,2,3})$. The essential local invariants are determined and interpreted geometrically. Several special classes of causal structures are considered including those that are a lift of pseudo-conformal structures and those referred to as causal structures with vanishing Wsf curvature. A rigidity theorem for causal structures of vanishing Wsf curvature over compact manifolds is obtained. A notion of half-flatness is defined for four dimensional causal structures whose null cones are ruled projective surfaces, which generalizes the notion of half-flat conformal structures of split signature. A twistorial construction for causal structures with vanishing Wsf curvature is given. Finally, the unique causal structure of split signature in dimension four with submaximal symmetry which does not descend to a pseudo-conformal structure is introduced.

CONTENTS

1. Introduction	1
1.1. Motivation and history	1
1.2. Main results	5
2. Local description of causal structures	7
2.1. Definitions and notational conventions	7
2.2. The flat model	9
2.3. The first order structure equations	10
2.4. Causal structures as parabolic geometries	21
2.5. Essential invariants and the Raychaudhuri equation	24
2.6. Examples	28
3. Causal structures with vanishing Wsf curvature	31
3.1. A compatibility condition	31
3.2. A rigidity theorem	33
3.3. Semi-integrable Lie contact structures	34
4. Four dimensions	37
4.1. Causal structures with ruled fibers	37
4.2. Submaximal causal structures of split signature	40
Appendix A. Full structure equations	41
References	47

1. INTRODUCTION

1.1. Motivation and history. In the General Theory of Relativity, space-time is represented by a four dimensional manifold M equipped with a Lorentzian metric g . An important feature of a Lorentzian structure is its associated field of null cones in TM given by the set of vectors $v \in TM$ satisfying $g(v, v) = 0$. As a result, it is possible to define a relation of *causality* between the points of M , i.e., a partial ordering $x < y$ defined

Date: April 11, 2017.

1991 *Mathematics Subject Classification.* 53B55 .

Key words and phrases. Causal geometry, Conformal geometry, Equivalence method, Cartan connection, Parabolic geometry.

by the property that x can be joined to y by a curve that is either time-like or null. If x and y represent two events in space-time then $x < y$ means that the occurrence of x has an effect on y . In [KP67] Kronheimer and Penrose examined causality in a manifold M of dimension n on an axiomatic basis. The starting point of their study is a continuous assignment of null cones in the tangent space of each point of M so that it is possible to define the set of points that are inside, outside and on the null cone in each tangent space.¹ The importance of the causal relationship between points of space-time became more significant in the context of the singularity theorems of Penrose and Hawking [Pen65, Pen72, HE73]. The study of causal relations in spaces equipped with a field of convex null cones in TM received a major contribution from the Alexandrov school in Russia under the name of *chronogeometry* [Ale67, Gut95], i.e., a geometry determined by the chronological relations between the points of the manifold. The objective was to use the principles of causality and symmetry to give an axiomatic treatment of special relativity. In [Seg76], Segal attempted to give such a treatment for general relativity (see also [DS01, Lev93, Wor84, Tau77, EPS72].) Subsequently, structures defined by assigning a convex cone in each tangent space, termed as causal structures, came to play a role in the Lie theory of semi-groups [HÓ97], causal boundaries [GPS05], and hyperbolic systems of partial differential equations [Sch13].

Rather than to investigate causality relations between the points of a manifold with a given causal structure, the main idea of this article is to gain a deeper understanding of the geometry of causal structures through the solution of the Cartan equivalence problem. This approach enables one to construct a complete set of local invariants and also to obtain some results on symmetries of causal structures. A global rigidity result can also be obtained through this approach.

Before going any further, a more precise definition of causal structures is in order. Given an $(n + 1)$ -dimensional manifold M , with $n \geq 3$, a *causal structure* (M, \mathcal{C}) of signature $(p + 1, q + 1)$ with $p, q \geq 0$, and $p + q = n - 1$ is given by a sub-bundle of the projectivized tangent bundle $\pi : \mathcal{C} \subset \mathbb{P}TM \rightarrow M$, whose fibers $\mathcal{C}_x \subset \mathbb{P}T_x M$ are projective hypersurfaces² with projective second fundamental form of signature (p, q) (see Definition 2.4). Note that in this definition the convexity of the null cones \mathcal{C}_x is not assumed.

Two causal structures (M, \mathcal{C}) and (M', \mathcal{C}') are locally equivalent around the points $p \in M$ and $p' \in M'$ if there exists a diffeomorphism $\psi : U \rightarrow U'$, where $p \in U \subset M$, $p' = \psi(p) \in U' \subset M'$, satisfying $\psi_*(\mathcal{C}_x) = \mathcal{C}_{\psi(x)}$ for all $x \in U$.

In order to describe a causal structure in an open set $U \subset TM$, one can use a defining function $L : TM \rightarrow \mathbb{R}$, $(x^i; y^j) \mapsto L(x^i; y^j)$, where (x^0, \dots, x^n) are coordinates for M and (y^0, \dots, y^n) are fiber coordinates. The function L is assumed to be homogeneous of some degree r_1 in the fiber variables and is referred to as a Lagrangian for the causal structure. Then, the vanishing set of L over U coincides with $U \cap \widehat{\mathcal{C}}$ where $\widehat{\mathcal{C}} \subset TM$ is the cone over $\mathcal{C} \subset \mathbb{P}TM$. The functions $L(x^i; y^i)$ and $S(x^i; y^i)L(x^i; y^i)$, where $S(x^i; y^i)$ is nowhere vanishing and homogeneous of degree r_2 in y^i 's, define the same causal structure.

As an example, if the fibers \mathcal{C}_x are hyperquadrics defined by a pseudo-Riemannian metric g of signature $(p + 1, q + 1)$, then the causal structure (M, \mathcal{C}) given by the family of null cones of g corresponds to the pseudo-conformal structure induced by the metric g .³ In other words, there exists a local defining function of this causal structure of the form $F(x^i; y^j) = g_{ij}(x)y^i y^j$.

The above description makes it clear that the relation of causal structures to pseudo-conformal structures is an analogue of what Finsler structures are to Riemannian structures. Recall that a Finsler structure (M^{n+1}, Σ^{2n+1}) is given by a codimension one sub-bundle $\Sigma \subset TM$, whose fibers $\Sigma_x \subset T_x M$ are strictly convex affine hypersurfaces, i.e., the second fundamental form of each fibers is positive definite. The sub-bundle Σ is called the indicatrix bundle (see [Bry02]).⁴

¹It should be said that in [KP67] it is assumed that each null cone is given by the vanishing locus of a quadratic form of signature $(n, 1)$.

²Here, a hypersurface is always taken to be a codimension one submanifold.

³A pseudo-conformal structure refers to the conformal class of a pseudo-Riemannian metric [AG96].

⁴Similarly, it is possible to define pseudo-Finsler structures of signature (p, q) , $p + q = n + 1$ by imposing the condition that the second fundamental form of the fibers Σ_x have signature $(p - 1, q)$.

The main motivation behind the study of causal structures comes from a variety of perspectives. As a part of the geometric study of differential equations, it was shown by Holland and Sparling [HS10] that there is locally a one-to-one correspondence between contact equivalence classes of third order ODEs $y''' = F(x, y, y', y'')$ and causal structures on their locally defined three dimensional solution space M . A causal structure is given in this setting by a family of curves $\pi : \mathcal{C} \rightarrow M$ where $\mathcal{C}_x := \pi^{-1}(x) \subset \mathbb{P}T_x M$ is a *non-degenerate* curve in the sense that it admits a well-defined projective Frenet frame. This causal structure descends to a conformal Lorentzian structure on M if a certain contact invariant of the third order ODE known as the Wünschmann invariant vanishes. In other words, it is shown in [HS10] that \mathcal{C} is locally isomorphic to $J^2(\mathbb{R}, \mathbb{R})$ ⁵ endowed with a foliation by lifts of contact curves on $J^1(\mathbb{R}, \mathbb{R})$ and that the Wünschmann invariant at each point of $J^2(\mathbb{R}, \mathbb{R})$ coincides with the projective curvature of the curve \mathcal{C}_x . As a result, the vanishing of the Wünschmann invariant implies that the curves \mathcal{C}_x are conics, i.e., the causal structure is conformal Lorentzian.

The work of Holland and Sparling is a part of the program of studying geometries arising from differential equations, which has a long history going back to Monge, Jacobi, Lie, Darboux, Goursat, Cartan and others. In the case of third order ODEs under contact transformations, Chern [Che40] used Cartan's method to solve the equivalence problem and showed that the solution depends on two essential relative invariants, namely the Wünschmann invariant $I(F)$ and the invariant $C(F) = \frac{\partial^4}{\partial q^4} F(x, y, p, q)$. Furthermore, he observed that if $I(F) = 0$, then the space of solutions of the ODE can be endowed with a conformal Lorentzian structure. Using Tanaka theory [Tan79], Sato and Yashikawa [SY98] showed how one can use Chern's $\{e\}$ -structure to associate a normal Cartan connection on some 10-dimensional principal bundle to the contact equivalence class of a third order ODE. In particular, they expressed the relative contact invariants of the ODE in terms of the curvature of the Cartan connection. Godlinski and Nurowski [GN09] explicitly computed the normal Cartan connection and explored other geometric structures that arise from a third order ODE. In particular, they used a result of Fox [Fox05b] to show if $C(F) = 0$, then the ODE induces a *contact projective structure*⁶ on $J^1(\mathbb{R}, \mathbb{R})$.

One can realize the works mentioned above on third order ODEs in terms of geometries arising from the double fibration

$$(1.1) \quad J^1(\mathbb{R}, \mathbb{R}) \xleftarrow{\tau} J^2(\mathbb{R}, \mathbb{R}) \xrightarrow{\pi} M$$

where the map π is the quotient map of $J^2(\mathbb{R}, \mathbb{R})$ by the foliation defined by the 2-jets of the solutions of the ODE. This map endows M with a causal structure. The null geodesics of M are given by $\pi \circ \tau^{-1}(p)$ where $p \in J^1(\mathbb{R}, \mathbb{R})$, while the contact geodesics are given by $\tau \circ \pi^{-1}(q)$ where $q \in M$.

Equivalently, given a causal structure (M, \mathcal{C}) in dimension three, Holland and Sparling showed that the double fibration above is equivalent to

$$\mathcal{N} \xleftarrow{\tau} \mathcal{C} \xrightarrow{\pi} M$$

where \mathcal{N} is the space of *characteristic curves* (or *null geodesics*)⁷ of the causal structure. The precise definition of these curves is given in Section 2.3.5. It turns out that starting with a causal structure (M, \mathcal{C}) , one can show that locally $\mathcal{C} \cong J^2(\mathbb{R}, \mathbb{R})$ and $\mathcal{N} \cong J^1(\mathbb{R}, \mathbb{R})$. Therefore, all the results mentioned above can be translated in terms of local invariants of the causal structure in three dimensions.

The correspondence above motivates the study of the local invariants of causal geometries in higher dimensions and differential equations that can be associated to them. As will be explained, this study turns out to be related to a number of current themes of research.

In [HS11], Holland and Sparling studied causal structures in higher dimensions and showed that an analogue of the Weyl sectional curvature can be defined. Also, they

⁵ In this article $J^k(\mathbb{R}^l, \mathbb{R}^m)$ denotes the space of k -jets of functions from \mathbb{R}^l to \mathbb{R}^m .

⁶ A contact path geometry over a contact manifold N is given by a family of contact paths with the property that at each point $x \in N$, a unique curve of the family passes through each contact direction in $T_x M$. A contact projective geometry is a contact path geometry whose contact paths are geodesics of some affine connection.

⁷ In this article the term characteristic curves is used to refer to the lift of null geodesics to the null cone bundle.

showed for causal structures the Raychaudhuri - Sachs equations from General Relativity remains valid for higher dimensional causal structures (also see [AJ16, Min15]).

Before describing the nature of the complete set of invariants for causal structures it is helpful to make parallels with what happens in Finsler geometry. On an $(n+1)$ -dimensional manifold M the indicatrix bundle Σ of a Finsler structure is $(2n+1)$ -dimensional equipped with a contact 1-form, i.e., a 1-form $\omega \in T^*\Sigma$ satisfying $\omega \wedge (d\omega)^n \neq 0$. The 1-form ω is called the Hilbert form of the Finsler structure. As indicated by Cartan [Car34, Car30] and Chern [Che48], in Finsler geometry there are two essential invariants comprising the total obstruction to local flatness, namely the *flag curvature* and the *centro-affine cubic form* of the fibers \mathcal{C}_x (also referred to as the Cartan torsion). The vanishing of the Cartan torsion implies that the Finsler structure is a Riemannian structure, in which case the flag curvature coincides with the sectional curvature.

Regarding causal structures, the main step is to realize that the $2n$ -dimensional space \mathcal{C} carries a quasi-contact structure (also known as even contact structure), i.e., a 1-form ω^0 satisfying $\omega^0 \wedge (d\omega^0)^{n-1} \neq 0$ and $(d\omega^0)^n = 0$. The 1-form ω^0 is referred to as the *projective Hilbert form*.⁸ It is shown that, for $n \geq 3$, there are two essential local invariants which form the total obstruction to local flatness. As one would expect, the *Fubini cubic form* of each fiber \mathcal{C}_x as a projective hypersurfaces in $\mathbb{P}T_x M$ constitutes one of these obstructions whose vanishing implies that the causal structure is a pseudo-conformal structure. The second essential invariant is a generalization of the sectional Weyl curvature of a pseudo-conformal structure restricted to the *shadow space*⁹ as found by Holland and Sparling in [HS11] (see Section 2.5). By analogy with Finsler geometry, the second essential invariant is called the *Weyl shadow flag curvature* (referred to as WSF curvature.)

Furthermore, from the first order structure equations for causal geometries it follows that a causal structure (M, \mathcal{C}) can be equivalently formulated as having a manifold \mathcal{C}^{2n} equipped with a distribution D of rank n which contains an involutive corank one sub-distribution $\Delta \subset D$ and satisfies a certain non-degeneracy condition. The (small) growth vector of D is $(n, 2n-1, 2n)$, i.e., $\text{rank}[D, D] = 2n-1$, and $\text{rank}[D, [D, D]] = 2n$.¹⁰ (see Section 2.4.2.) In this case, M^{n+1} is the space of integral leaves of Δ .¹¹ This description identifies causal geometries as parabolic geometries modeled on $B_n/P_{1,2}$ and $D_n/P_{1,2}$ for $n \geq 4$ and $D_3/P_{1,2,3}$, with their natural $[3]$ -grading. As will be shown, they have some similar features to path geometries [Gro00] and contact path geometries [Fox05a], which correspond respectively to parabolic geometries modeled on $A_n/P_{1,2}$ and $C_n/P_{1,2}$.

Another essential part of any causal structure is given by its set of *characteristic curves*. Note that as a quasi-contact manifold, \mathcal{C} is equipped with the characteristic line field of the projective Hilbert form, i.e., the unique degenerate direction of $d\omega^0$ lying in $\text{Ker } \omega^0$, defined by vector fields \mathbf{v} satisfying

$$(1.2) \quad \omega^0(\mathbf{v}) = 0, \quad \mathbf{v} \lrcorner d\omega^0 = 0.$$

The integral curves of a characteristic vector field \mathbf{v} is called a characteristic curve. In the case of pseudo-conformal structures the projection of these curves to M coincide with null geodesics.

The description above shows that, unlike Finsler geometry, the definition of characteristic curves does not depend on a choice of metric. Recall that in Finsler geometry geodesics can be defined as the extremals of the arc-length functional

$$\Phi(\gamma) = \int_I \gamma^* \phi,$$

where $\gamma : (a, b) \rightarrow \mathcal{C}$ is a curve in \mathcal{C} and $\gamma^* \phi = L(x^i; y^i) dt$. The function $L(x^i; y^i)$ is called the Finsler metric which in the calculus of variations is referred to as the Lagrangian.

⁸This terminology is introduced in analogy with the Hilbert form in Finsler geometry and the calculus of variations [BCS00].

⁹The terminology is due to a physical interpretation made by Sachs [Sac61] according to which if an object follows a congruence of null geodesics, its infinitesimal shadow casts onto the pull-back of this space. This term was initially used by Holland and Sparling in [HS11].

¹⁰The sections of the distribution $[D, D]$ are given by $\Gamma([D, D]) = [\Gamma(D), \Gamma(D)]$.

¹¹In [ANN15], these structures are referred to as the B_n and D_n classes of what they call parabolic geometries of Monge type.

The Finsler metric is homogeneous in y^i 's and has non-degenerate vertical Hessian¹². When addressing the geodesics of a Finsler structure, or more generally extremal manifolds in the calculus of variations, it is generally assumed that the Lagrangian does not vanish along them [Gri83]. The study of extremal curves along which the Lagrangian vanishes seems to have received much less attention in the literature (see [Run83, Run84]). The exception is the case of Lagrangians that are quadratic in y^i 's. This is due to the fact that they arise in the context of pseudo-conformal structures. For instance, this problem is briefly discussed by Guillemin [Gui89], where he studies the deformations of the product metric $\pi_1^* dt^2 \oplus -\pi_2^* g$ on $S^1 \times S^2$ where π_1 and π_2 are projections onto S^1 , and S^2 and g is the round metric on S^2 , which is *Zollfrei*, i.e., all of its null geodesics are closed with the same period. In Section 2.5.3 it is discussed how causal structures provide a geometric setup for the variational problems involving certain classes of Lagrangians $L(x^i; y^i)$ which are homogeneous in y^i 's and have vertical Hessian of maximal rank over their vanishing set.

The foliation of \mathcal{C} defined by the characteristic curves allows one to define the quotient map $\tau : \mathcal{C} \rightarrow \mathcal{N}$ where \mathcal{N} is $(2n - 1)$ -dimensional and is called the space of characteristic curves of \mathcal{C} . A priori, \mathcal{N} is only locally defined around generic points and can be used to give a geometric description of the theory of correspondence spaces for causal structure in the sense of Čap [Čap05] (see Section 3.3). Note that the space \mathcal{N} , when globally defined, is of interest in Penrose's twistor theory [LeB90] and a twistorial description of causal relations among the points of the space-time [Low06].

Finally, an interesting instance of studying causal structure can be found in [Hwa13] where Hwang considers complex causal structures arising on certain uniruled projective manifolds, M^{n+1} , satisfying the property that through a generic point $x \in M$ there passes an $(n - 1)$ -parameter family of *rational curves of minimal degree*. The hypersurface $\mathcal{C}_x \subset \mathbb{P}T_x M$ obtained from the tangent directions to such curves at the point x is called the variety of minimal rational tangent (or VMRT). Under the assumption that \mathcal{C}_x is smooth of degree ≥ 3 , he shows that the causal structure defined by them is locally isotrivially flat (see Definition 2.18)¹³. Furthermore, it turns out that the characteristic curves for such causal structures coincide with rational curves of minimal degree¹⁴. Finally, assuming that M has Picard number 1 and $n \geq 2$, he shows that M is biregular to a hyperquadric equipped with its natural causal structure (see Section 2.2). This theorem is a generalization of what has been considered previously in [Bel01, Ye94] for complex conformal structures.

More broadly, Hwang's study of causal structures is a part of the differential geometric characterization of uniruled varieties initiated by Hwang and Mok [Hwa12] via geometric structures referred to as *cone structures* (see Definition 2.1). The starting point for Hwang and Mok is a given family of rational curves of minimal degree whose tangent directions at a generic point $x \in M$ give rise to the VMRT $\mathcal{C}_x \subset \mathbb{P}T_x M$. For instance, if $\mathcal{C}_x = \mathbb{P}T_x M$ one obtains the complex analogue of path geometry [Gro00] and if \mathcal{C}_x is a hyperplane which induces a contact distribution on M , one obtains a complex contact path geometry [Fox05a]. The program of Hwang and Mok seems interesting and calls for further exploration from the perspective of the method of equivalence, especially through the lens of flag structures as developed in [DZ12].

1.2. Main results. In all the statements of this article smoothness of underlying manifolds and maps is always assumed and the dimension of M is always taken to be at least four unless otherwise specified.

In Section 2, after reviewing the flat model for causal structures, the first order structure equations for a causal geometry (M^{n+1}, \mathcal{C}) of signature $(p+1, q+1)$ are derived. It is shown that the bundle of null cones \mathcal{C} is foliated by characteristic curves and is endowed with the conformal class of a quadratic form which is degenerate along the vertical directions

¹²It is usually assumed that the Finsler metric when restricted to $T_x M$ only vanishes at the origin and is positive elsewhere [BCS00]. However, as was mentioned previously, the condition can be relaxed [She01].

¹³Hwang refers to such structures as flat cone structures of codimension one.

¹⁴Motivated by the notion of *conformal torsion* introduced by LeBrun [LeB83] and *contact torsion* introduced by Fox [Fox05a], one can define *causal torsion* whose vanishing implies that the rational curves of minimal degree coincide with the characteristic curves of the causal structure they define. The precise definition of causal torsion is not given in this article.

of the fibration $\mathcal{C} \rightarrow M$. Subsequently, the first order structure equations are used in the Appendix to obtain structure equations leading to an $\{e\}$ -structure.

The essential theorem of this article that is stated in Section 2.3.13 is the following.

Theorem 2.12. *To a causal structure $(M^{n+1}, \mathcal{C}^{2n})$, one can associate an $\{e\}$ -structure on a principal bundle $P_{1,2} \rightarrow \mathcal{P} \rightarrow \mathcal{C}$ of dimension $\frac{(n+3)(n+2)}{2}$ where $P_{1,2} \subset O(p+2, q+2)$ is the parabolic subgroup defined in (2.5). The essential invariants of a causal structure are the Fubini cubic form of the fibers and the WSF curvature. The vanishing of the essential invariants implies that the $\{e\}$ -structure coincides with the one defined by the Maurer-Cartan forms of $\mathfrak{o}(p+2, q+2)$.*

The proof of the above theorem takes up most of the Appendix. Using the $\{e\}$ -structure thus obtained, it is straightforward to deduce that the essential invariants are the Fubini cubic form and the WSF curvature. Section 2.4 deals with the parabolic nature of causal structures. In Section 2.5 some geometrical interpretations of the essential invariants are given. The first Section ends with examples of causal structures.

In Section 3.1, the structure equations for causal geometries with vanishing WSF curvature are examined. It is shown that if the Lie derivative of the Fubini cubic form along the vector fields tangent to the characteristic curves is proportional to itself, then the WSF curvature has to vanish, unless the causal geometry is a pseudo-conformal structure. Such spaces are the causal analogues of Landsberg spaces in Finsler geometry¹⁵. Moreover, having defined the scalar shadow flag curvature (referred to as the SSF curvature), and the notion of completeness of characteristic curves for a causal structure, the following theorem is proved in Section 3.2.

Theorem 3.5. *Let (M, \mathcal{C}) be a causal structure with vanishing WSF curvature over a compact manifold with non-zero Fubini cubic form. If the characteristic curves of \mathcal{C} are complete, then the scalar flag curvature of (M, \mathcal{C}) is non-negative.*

The above theorem implies the following corollary.

Corollary 3.6. *Let (M, \mathcal{C}) be a compact causal structure with vanishing WSF curvature and complete characteristic curves. If it has negative SSF curvature, then it is a flat pseudo-conformal structure.*

The corollary above can be taken as a rigidity theorem which is a causal analogue of the seminal work of Akbar-Zadeh in Finsler geometry [AZ88].

Section 3.3 involves the induced structure on the space of characteristic curves, \mathcal{N} , of a causal structure with vanishing WSF curvature. The $(2n-1)$ -dimensional manifold \mathcal{N} is shown to have a contact structure and is endowed with an $(n+1)$ -parameter family of Legendrian submanifolds with the property that through each point $\gamma \in \mathcal{N}$ there passes a 1-parameter family of them. Moreover, the sub-bundle of $T\mathcal{N}$ defined by the contact distribution contains a Segré cone of type $(2, n-1)$. The theorem below is proved in which the property of being *semi-integrable* means that at every point $\gamma \in \mathcal{N}$, the Segré cone is ruled by the tangent spaces of the corresponding 1-parameter family of Legendrian submanifolds passing through γ .

Theorem 3.10. *Given a causal structure with vanishing WSF curvature (\mathcal{C}, M) , its space of characteristic curves admits a semi-integrable Lie contact structure. Conversely, any semi-integrable Lie contact structure on a manifold \mathcal{N} induces a causal structure with vanishing WSF curvature on the space of its corresponding Legendrian submanifolds.*

It should be noted that the theorem above is reminiscent of Grossman's work [Gro00] on torsion-free path geometries and the term semi-integrability is borrowed from there. He showed that given a torsion-free path structure on N^{n+1} , the space of paths, \mathcal{S}^{2n} , has a Segré structure, i.e., each tangent space is equipped with a Segré cone of type $(2, n)$. Furthermore, \mathcal{S} contains an $(n+1)$ -parameter family of n -dimensional submanifolds such that through each point $\gamma \in \mathcal{S}$ there passes a 1-parameter family of them. Grossman showed that at each point $\gamma \in \mathcal{S}$, the Segré cone is ruled by the tangent spaces of the 1-parameter family of submanifolds passing through γ . He called such Segré structures *semi-integrable*. Furthermore, twistorial constructions such as that of Theorem 3.10 fit exactly into Čap's

¹⁵See [BCS00] for an account.

theorem of correspondence spaces [Čap05]. However, Čap's theory does not contain the geometric interpretation given here in terms of existence of submanifolds that rule the null cone or the Segré cone. Such interpretations are clear from the constructions presented here.

In Section 4.1 a notion of half-flat causal structures of signature $(2, 2)$ is introduced using the existence of a 3-parameter family of surfaces whose tangent planes at each point of M give a ruling of the projective surfaces \mathcal{C}_x . This notion of half-flatness coincides with the notion of half-flatness for conformal structures of signature $(2, 2)$.

Finally, in Section 4.2 a study of the symmetry algebra of certain non-flat causal structures of split signature that do not induce a conformal structure on M is given from which the proposition below follows.

Proposition 4.8. *The submaximal causal structure of split signature that does not descend to a pseudo-conformal structure has 8 dimensional symmetry algebra and is locally equivalent to the locally isotrivially flat causal structure whose null cones are projectively equivalent to Cayley's cubic scroll.*

Locally isotrivially flat causal structures, defined in 2.18, are the causal analogue of locally Minkowskian Finsler spaces.

The Appendix contains the details of the equivalence problem calculations. Since causal structures are examples of parabolic geometries of type $(B_n, P_{1,2})$ and $(D_n, P_{1,2})$ if $n \geq 4$ and $(D_3, P_{1,2,3})$, they admit a *regular* and *normal* Cartan connection ([ANN15, ČS09]). Using the machinery of Tanaka, the correct change of coframe can be given recursively, in order to introduce a Cartan connection on the structure bundle of causal geometries which is normal in the sense of Tanaka. However, for computational reasons, writing down the precise form of the Cartan connection leads to very long expressions which are not particularly enlightening (see [Mak16]). Furthermore, the first order structure equations suffice for the purposes of this article.

Acknowledgments. This work was carried out under the guidance of Professor N. Kamran, whom I would like to thank greatly for his clear explanation of Cartan's method of equivalence and introducing me to the work of Holland and Sparling [HS10]. His encouragement, help and support during my PhD made this work possible. I am also very much indebted to Dennis The for the many helpful conversations that we had and for patiently explaining some aspects of parabolic geometry. The funding for this work was provided by McGill University.

2. LOCAL DESCRIPTION OF CAUSAL STRUCTURES

In this section, after introducing the necessary definitions and notations, the first order structure equations of causal structures are derived. It is assumed that the reader is familiar with Cartan's method of equivalence for G -structures (see [Gar89, Olv95] for an account). The essential torsion terms appearing in the first structure equations are interpreted geometrically. The derivation of the full structure equations is carried out in the Appendix A. In Section 2.5 a discussion on the fundamental invariants of a causal structure is given. Finally, the section ends with a few examples of causal structures.

2.1. Definitions and notational conventions. Since the main purpose of this article is the local study of causal geometries, the smoothness of manifolds and maps that are considered is always assumed. In other words, the manifolds considered are assumed to be the maximal open sets over which the necessary smoothness conditions are satisfied.

Let M be a smooth $(n + 1)$ -dimensional (real or complex) manifold, $n \geq 3$ and $\mathbb{P}TM$ denote its projectivized tangent bundle with projection $\pi : \mathbb{P}TM \rightarrow M$, and fibers $\pi^{-1}(x) = \mathbb{P}T_xM$. Given a tangent vector $y \in T_xM$, its projective equivalence class is denoted by $[y] \in \mathbb{P}T_xM$. For $S \subset \mathbb{P}T_xM$, the cone over S is defined as

$$\widehat{S} := \{y \in T_xM \mid [y] \in S\} \subset T_xM.$$

Throughout this article, the indices range as

$$0 \leq i, j, k \leq n, \quad 1 \leq a, b, c \leq n - 1, \quad 0 \leq \alpha, \beta \leq n - 1,$$

and are subjected to the summation convention. When a set of 1-forms is introduced as $\mathcal{I} = I\{\omega^i\}_{i=0}^n$, it is understood that \mathcal{I} is the ideal algebraically generated by $\{\omega^0, \dots, \omega^n\}$ in the exterior algebra of M .

The symmetrization and anti-symmetrization operations for tensors are denoted using Penrose's abstract index notation. For example, consider a tensor with coefficients A_{abcd} . The tensor $A_{(ab|c|d)}$ is symmetric with respect to all indices except the third, and $A_{[a|bc|d]}$ is anti-symmetric in the first and last indices. When dealing with symmetric forms such as $g = 2\omega^0 \circ \omega^n - \varepsilon_{ab}\omega^a \circ \omega^b$, or $F_{abc}\theta^a \circ \theta^b \circ \theta^c$ with $F_{abc} = F_{(abc)}$, the symbol \circ denotes the symmetric tensor product. Bold-face letters such as $\mathbf{i}, \mathbf{j}, \mathbf{a}, \mathbf{b}$ are used to denote a specific index, e.g. the set $\{\omega^{\mathbf{a}}\}$ only contains the 1-form $\omega^{\mathbf{a}}$. The summation convention is not applied to bold-face letters.

At this stage, before stating the definition of a causal structure, a more general geometric structure, namely that of a *cone structure*, is defined [Hwa12]. The reason for this is that in the process of coframe adaption for cone structures of codimension one, causal structures are distinguished as a sub-class of such geometries via the characterizing property of having *tangentially non-degenerate* fibers (see Section 2.3.3).

Definition 2.1. A *cone structure* of codimension $k \leq n$ on M^{n+1} is given by an immersion $\iota : \mathcal{C} \rightarrow \mathbb{P}TM$ where \mathcal{C} is a connected, smooth manifold of dimension $2n + 1 - k$ with the property that the fibration $\pi \circ \iota : \mathcal{C} \rightarrow M$ is a submersion whose fibers $\mathcal{C}_x := (\pi \circ \iota)^{-1}(x)$ are mapped, via the immersion $\iota_x : \mathcal{C}_x \rightarrow \mathbb{P}T_x M$, to connected projective submanifolds of codimension k in $\mathbb{P}T_x M$.

Two cone structures $\iota : \mathcal{C} \rightarrow \mathbb{P}TM$ and $\iota' : \mathcal{C}' \rightarrow \mathbb{P}TM'$ are locally equivalent around points $x \in M, x' \in M'$, if there exists a diffeomorphism $\phi : U \rightarrow U'$ where $x \in U \subset M$ and $x' \in U' \subset M'$, such that $x' = \phi(x)$ and $\phi_*(\iota(\mathcal{C})) = \iota'(\mathcal{C}'_{\phi(y)})$ for all $y \in M$.

In this article, only cone structures of codimension one are considered i.e., the fibers $\iota(\mathcal{C}_x) \subset \mathbb{P}T_x M$ are projective hypersurfaces. Since the map $\iota : \mathcal{C} \rightarrow \mathbb{P}TM$ is assumed to be an immersion and this article concerns the local nature of geometric structures, it is safe to drop the symbol ι in $\iota(\mathcal{C})$ and $\iota(\mathcal{C}_x)$ and denote them as \mathcal{C} and \mathcal{C}_x , respectively. As a result, by restricting to small enough neighborhoods, one can safely assume $\mathcal{C} \subset \mathbb{P}TM$ is a sub-bundle defined in an open set $U \subset \mathbb{P}TM$, where the fibers \mathcal{C}_x are smooth submanifolds. Consequently, a cone structure of codimension one over a manifold M can be represented by (\mathcal{C}, M) .

In local coordinates, a point of TM is represented by $(x; y)$ where $x = (x^0, \dots, x^n)$ are the base coordinates and $y = (y^0, \dots, y^n)$ are the fiber coordinates. In a sufficiently small neighborhood of a point $p = (x; z) \in \mathcal{C}$, the homogeneous coordinates $[y^0 : \dots : y^n]$ can be chosen so that $z = [0 : \dots : 0 : 1]$ and the cone $\widehat{\mathcal{C}}_x$ is tangent to the hyperplane $\{y_0 = 0\}$ along \hat{z} . The affine coordinates (y^0, \dots, y^{n-1}) obtained from setting $y^n = 1$, are used to derive the explicit expressions of the invariants. Using these preferred coordinates in a sufficiently small neighborhood of p , it is assumed that \mathcal{C} is given in terms of homogeneous coordinates as a locus

$$(2.1) \quad L(x^0, \dots, x^n; y^0, \dots, y^n) = 0$$

where L is homogeneous of degree r in y^i and, in terms of affine coordinates, as a graph

$$(2.2) \quad y^0 = F(x^0, \dots, x^n; y^1, \dots, y^{n-1}).$$

Example 2.2. A cone structure of codimension one on M^{n+1} whose fibers are hyperplanes given by $\text{Ker } \omega$ for some $\omega \in T^*M$, and are *maximally non-integrable*, i.e., $\omega \wedge (d\omega)^{k-1} \neq 0$, is called a *contact structure* if $n = 2k - 1$, and a *quasi-contact structure* (or *even contact structure*) if $n = 2k$. These can be considered as “degenerate” examples of codimension one cone structures and have no local invariants. In these cases, the function F in (2.2) is of the form $F(x; y) = c_a(x)y^a + c_n(x)$, with $\omega = dx^0 + c_a(x)dx^a + c_n(x)dx^n$.

Example 2.3. A pseudo-conformal structure, i.e., a conformal class of a pseudo-Riemannian metric g_{ij} of signature $(p + 1, q + 1)$, $p, q \geq 0$, can be introduced by its field of null cones, i.e., the set of quadrics

$$\mathcal{C}_x := \{[y] \in \mathbb{P}T_x \mid g_{ij}y^i y^j = 0\},$$

As a result, a pseudo-conformal structure is an example of a cone structure of codimension one, which, in contrast to the previous example, is “non-degenerate” (see section 2.3.3).

In terms of the local form (2.2), a representative g_{ij} of the conformal class, at a point $s = (x; y) \in \mathcal{C}$, can be put in the form $2y^0 y^n - \varepsilon_{ab} y^a y^b$, where

$$(2.3) \quad [\varepsilon_{ab}] = \mathbb{I}_{p,q} := \begin{pmatrix} \mathbb{I}_p & 0 \\ 0 & -\mathbb{I}_q \end{pmatrix},$$

and therefore $F(s) = \varepsilon_{ab} y^a y^b$.

In Section 2.3.3, when a coframe $(\omega^0, \dots, \omega^n, \theta^1, \dots, \theta^{n-1})$ over \mathcal{C} is introduced, the corresponding coframe derivatives are denoted by

$$dG = G_{;i} \omega^i + G_{;\underline{a}} \theta^{\underline{a}}.$$

2.2. The flat model. When setting up the equivalence problem for a given geometric structure, finding the flat model can be very helpful for identifying the structure bundle. Naturally, one expects the flat model for causal structures to coincide with the flat model for conformal geometry. In causal geometry, the flat model is obtained from the natural pseudo-conformal structure on a quadric. Consider the quadric $Q_{p+1,q+1}^{n+1} \subset \mathbb{P}^{n+2}$, where $p+q = n-1$, given by ${}^t v S v = 0$, where $v = {}^t(y, x^0, x^1, \dots, x^n, z) \in \mathbb{R}^{n+3}$ and

$$S := \begin{pmatrix} 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & \mathbb{I}_{p,q} & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 \end{pmatrix}$$

has signature $(p+2, q+2)$. It is equipped with a pseudo-conformal structure of signature $(p+1, q+1)$, expressed as $2y^0 y^n - \varepsilon_{ab} y^a y^b$ where y^i 's denote the fiber coordinates of $TQ_{p+1,q+1}^{n+1}$. The null cone bundle of this pseudo-conformal class is denoted by $\mathcal{C} \subset \mathbb{P}TQ_{p+1,q+1}^{n+1}$. Consider the *flag manifold*

$$(2.4) \quad \mathbb{NF}_{0,1}^{n+1} := \{(V_1, V_2) \in \text{Gr}_1^0(n+3) \times \text{Gr}_2^0(n+3) : V_1 \subset V_2\}$$

where $\text{Gr}_k^0(m)$ denotes the space of k -dimensional subspaces of \mathbb{R}^m which are null with respect to the inner product S , e.g., $\text{Gr}_1^0(n+3) \cong Q_{p+1,q+1}^{n+1}$. Taking a basis $\{e_1, \dots, e_m\}$ for \mathbb{R}^m , the transitivity of the action of $G_{p,q} := O(p+2, q+2)$ on $\text{Gr}_k^0(m)$ results in isomorphisms $\text{Gr}_k^0(m) \cong G_{p,q}/P_k$ where P_k is the parabolic subgroup isomorphic to the stabilizer of the subspace spanned by $\{e_1, \dots, e_k\}$.

Let $m = n+3$, $n \geq 4$ and define the parabolic subgroup $P_{k_1, \dots, k_i} := P_{k_1} \cap \dots \cap P_{k_i}$ of G . Consequently, it follows that

$$(2.5) \quad \begin{aligned} \mathbb{NF}_{0,1}^4 &\cong G_{p,q}/P_{1,2,3}, \quad p+q=2, \\ \mathbb{NF}_{0,1}^{n+1} &\cong G_{p,q}/P_{1,2}, \quad p+q=n-1 \geq 3 \end{aligned}$$

The relevant fibrations are shown in the diagram below.

$$(2.6) \quad \begin{array}{ccccc} & & G_{p,q} & & \\ & & \downarrow v & & \\ Q_{p+1,q+1}^{n+1} & \xleftarrow{\pi} & \mathbb{NF}_{0,1}^{n+1} & \xrightarrow{\tau} & \text{Gr}_2^0(n+3) \end{array}$$

Note that $\mathbb{NF}_{0,1}^{n+1}$ is isomorphic to \mathcal{C} . The Maurer-Cartan form on $G_{p,q}$ determines a canonical Cartan connection over \mathcal{C} which, as will be shown, is associated to the flat causal structure. Moreover, the fibration of $G_{p,q}$ over $\text{Gr}_2^0(n+3)$ gives the flat model for Lie contact structures studied in [SY89, Miy91]. As will be seen, these structures arise on the space of characteristic curves of certain classes of causal geometries.

The Maurer-Cartan form of $G_{p,q}$ can be written as

$$(2.7) \quad \varphi = \begin{pmatrix} \phi_0 + \phi_n & -\pi_n & -\pi_b & -\pi_0 & 0 \\ \omega^n & \phi_0 - \phi_n & -\gamma_b & 0 & -\pi_0 \\ \omega^a & -\theta^a & -\phi^a_b & -\gamma^a & \pi^a \\ \omega^0 & 0 & -\theta_b & -\phi_0 + \phi_n & -\pi_n \\ 0 & \omega^0 & -\omega_b & \omega^n & -\phi_0 - \phi_n \end{pmatrix},$$

with $d\varphi + \varphi \wedge \varphi = 0$. The symbol ε_{ab} is used to lower and raise indices. The set of semi-basic 1-forms with respect to the projections $v, v \circ \pi$ and $v \circ \tau$ are respectively

$$\begin{aligned}\mathcal{I}_1 &= \{\omega^0, \dots, \omega^n, \theta^1, \dots, \theta^{n-1}\}, \\ \mathcal{I}_2 &= \{\omega^0, \dots, \omega^n\}, \\ \mathcal{I}_3 &= \{\omega^0, \dots, \omega^{n-1}, \theta^1, \dots, \theta^{n-1}\}.\end{aligned}$$

The tangent spaces $T_p\mathcal{C}$ are equipped with a filtration defined as follows. At $p \in \mathcal{C}$, the line over p , denote by $\hat{p} \in T_{\pi(p)}Q_{p+1,q+1}^{n+1}$, is null. Let ω^0 denote a representative of the dual projective class of \hat{p} with respect to the conformally defined inner product S . For instance, if $p = [e_n]$, then take $\omega^0 = e_n^\flat$. Define $K^2 := \text{Ann}(\pi^*\omega^0) \subset T\mathcal{C}$. Using the tautological bundle

$$K^1 := \{v \in T_p\mathcal{C} : (\pi_*(p))(v) \in \hat{p} \subset T_{\pi(p)}Q_{p+1,q+1}^{n+1}\},$$

the filtration $K^1 \subset K^2 \subset T\mathcal{C}$ is obtained. At $p \in \mathcal{C}$, let $K_{-1}(p) := K^1(p)$, $K_{-2} := K^2(p)/K^1(p)$ and $K_{-3} := K^3(p)/K^2(p)$. The graded vector space

$$(2.8) \quad \mathfrak{m}(p) = K_{-1}(p) \oplus K_{-2}(p) \oplus K_{-3}(p)$$

can be associated to the filtration of $T_p\mathcal{C}$. Note that at each $p \in \mathcal{C}$, $\mathfrak{m}(p) \cong \mathfrak{m}$. The Lie bracket on $T\mathcal{C}$ induces a bracket on \mathfrak{m} which is tensorial and makes \mathfrak{m} a graded nilpotent Lie algebra (GNLA), i.e., $[K_i, K_j] = K_{i+j}$ with $K_i = 0$ for all $i \geq 4$ and K_{-1} the generating component. Let G_0 denote the subgroup of automorphisms of $T\mathcal{C}$, i.e., $O(p+2, q+2)$, which preserves the grading of \mathfrak{m} via the adjoint action. Let K_0 denote the Lie algebra of G_0 . Then, the Lie algebra of automorphisms of \mathcal{C} , i.e., $\mathfrak{g} = \mathfrak{o}(p+2, q+2)$, admits a \mathbb{Z} -grading defined by

$$\mathfrak{g}_k := \{g \in \mathfrak{g} : g \cdot K_i \subset K_{i+k} \text{ for } 0 \geq i \geq -3\}.$$

This is a $|\mathbb{Z}|$ -grading, i.e., $\mathfrak{g} = \bigoplus_{i=-3}^3 \mathfrak{g}_i$ such that $[\mathfrak{g}_i, \mathfrak{g}_j] \subset \mathfrak{g}_{i+j}$ with $\mathfrak{g}_i = 0$ for $|i| > 3$, and \mathfrak{g}_{-1} generates $\mathfrak{g}_- := \mathfrak{g}_{-1} \oplus \mathfrak{g}_{-2} \oplus \mathfrak{g}_{-3}$.

Note that $K^1 = E \oplus \text{Ver}$ where Ver is the vertical distribution with respect to the fibration π and $E = \pi_*(\mathfrak{g}_{-1})$. Moreover, $T\mathcal{C}$ is equipped with a conformally defined quadratic form obtained by lifting the canonical one defined on $Q_{p+1,q+1}^{n+1}$,

$$g = 2\omega^0 \circ \omega^n - \varepsilon_{ab}\omega^a \circ \omega^b,$$

which is degenerate on Ver . The group of automorphisms of $T\mathcal{C}$, which preserves the grading of \mathfrak{m} , also preserves the conformal class of the lifted quadratic form g on $T\mathcal{C}$. The Lie algebra $\mathfrak{g}_0 \subset \mathfrak{g}$ which corresponding to the group of automorphisms of \mathfrak{m} that preserves the grading, is isomorphic to $\mathfrak{co}(p, q) \oplus \mathbb{R}$.

Using the Maurer-Cartan form φ , the subspaces $\mathfrak{g}_{-3}, \mathfrak{g}_{-2}$, and \mathfrak{g}_{-1} are dual to the Pfaffian systems $\mathcal{I}_{-3} := \{\omega^0\}$, $\mathcal{I}_{-2} := \{\omega^a\}_{a=1}^{n-1}$, and $\mathcal{I}_{-1} := \{\omega^n, \theta^a\}_{a=1}^{n-1}$.

2.3. The first order structure equations. In this section the first order structure equations for causal geometries are derived via the method of equivalence. The procedure involves successive coframe adaptations, as a result of which various important aspects of causal geometries are unraveled, e.g., the quasi-contact structure of \mathcal{C} , the projective invariants of the fibers \mathcal{C}_x , and the WSF curvature. Also, it is noted that a causal structure can be equivalently formulated as a codimension one sub-bundle of $\mathbb{P}T^*M$, which is called a dual formulation of causal geometries. Such formulation is traditionally preferred in the context of geometric control theory [AZ02].

The derivation of the first order structure equations is similar to that of Finsler structures as presented in [Bry02].

2.3.1. The projective Hilbert form. Given a cone structure of codimension one (\mathcal{C}, M) , the subset $\mathcal{C}_x \subset \mathbb{P}T_xM$ is a projective hypersurface, for all $x \in M$. At a point $y \in \mathcal{C}_x$, the affine tangent space $\widehat{T}_y\mathcal{C}_x$ is defined as the hyperplane $T_z\widehat{\mathcal{C}}_x \subset T_xM$ tangent to the cone $\widehat{\mathcal{C}}_x \subset T_xM$, containing $z \in \widehat{y}$. Note that this subspace is independent of the choice of z . It is important here to distinguish between the affine tangent space $\widehat{T}_y\mathcal{C}_x \subset T_xM$ and the vertical tangent space $VT_{(x;y)}\mathcal{C}$ corresponding to the fibration $\pi \circ \iota : \mathcal{C} \rightarrow M$.

The 1-form $\omega^0 \in T^*\mathcal{C}$, at $(x; y) \in \mathcal{C}$ is defined as a multiple of $(\pi \circ \iota)^*(\alpha)$ where $\alpha := \text{Ann}(\widehat{T}_y\mathcal{C}_x) \in T_x^*M$. By definition, ω^0 is defined up to a multiplication by a nowhere vanishing smooth function on \mathcal{C} and is therefore semi-basic with respect to the fibration $\pi : \mathcal{C} \rightarrow M$. In this article ω^0 is referred to as the projective Hilbert form.

2.3.2. *The adapted flag.* At each point $(x; y) \in \mathcal{C}$, there is a natural basis $(\frac{\partial}{\partial \omega^0}, \dots, \frac{\partial}{\partial \omega^n})$ for $T_x M$ adapted to the flag

$$\hat{y} \subset \hat{T}_y \mathcal{C}_x \subset T_x M,$$

where $\{\frac{\partial}{\partial \omega^n}\}$ and $\{\frac{\partial}{\partial \omega^1}, \dots, \frac{\partial}{\partial \omega^n}\}$, respectively, span \hat{y} and $\hat{T}_y \mathcal{C}_x$.

In terms of the graph (2.2), the local expressions are

$$(2.9) \quad \begin{aligned} \frac{\partial}{\partial \omega^n} &= F \frac{\partial}{\partial x^0} + y^a \frac{\partial}{\partial x^a} + \frac{\partial}{\partial x^n}, \\ \frac{\partial}{\partial \omega^a} &= \partial_a F \frac{\partial}{\partial x^0} + \frac{\partial}{\partial x^a} \\ \frac{\partial}{\partial \omega^0} &= \frac{\partial}{\partial x^0}. \end{aligned}$$

The dual coframe $(\omega^0, \dots, \omega^n)$ can be written as

$$(2.10) \quad \begin{aligned} \omega^n &= dx^n, \\ \omega^a &= dx^a - y^a dx^n \\ \omega^0 &= dx^0 - \partial_a F dx^a + (y^a \partial_a F - F) dx^n. \end{aligned}$$

The pull-back of these 1-forms to \mathcal{C} spans the semi-basic 1-forms with respect to the fibration $\pi : \mathcal{C} \rightarrow M$. At each point $p \in \mathcal{C}$, given by $p = (x^i; y^i)$, these 1-forms are adapted to the flag

$$(2.11) \quad VT_p \mathcal{C} \subset K_p \subset H_p \subset T_p \mathcal{C},$$

where $VT_p M = \text{Ker} \{\omega^0, \dots, \omega^n\}$ is the vertical tangent bundle, $K_p = \text{Ker} \{\omega^0, \dots, \omega^{n-1}\}$ is the fiber of the tautological vector bundle given by

$$K_p = \{v \in T_p \mathcal{C} \mid \pi_*(p)(v) \in \hat{y} \subset T_{\pi(p)} M\},$$

where $p = (x; y)$, the line \hat{y} corresponds to the projective class of y and $H_p = \text{Ker} \{\omega^0\}$ is the quasi-contact distribution (see Section 2.3.5). To clarify the notation, if $p = (x; y)$ then the fiber $VT_p \mathcal{C}$ is also denoted by $T_y \mathcal{C}_x$ and the quasi-contact distribution can be expressed as the splitting $T_y \mathcal{C}_x \oplus \hat{T}_y \mathcal{C}_x$ which is not unique.

2.3.3. *Causal structures and the first coframe adaptation.* Recall that a projective hypersurface $\mathcal{V} \subset \mathbb{P}^n$ is called *tangentially non-degenerate* if its projective second fundamental form has maximal rank (see [Sas99, AG93]). Now the stage is set to define causal structures.

Definition 2.4. A *causal structure* of signature $(p+1, q+1)$ on M^{n+1} is a cone structure of codimension one, (M, \mathcal{C}) , where the fibers \mathcal{C}_x are tangentially non-degenerate projective hypersurfaces and their projective second fundamental form has signature (p, q) everywhere.

In section 2.3.6 it is shown that the tangential non-degeneracy assumption of the fibers \mathcal{C}_x implies that the signature of h^{ab} at any point of \mathcal{C} determines its the signature of h^{ab} everywhere.

Let d_V denote the vertical exterior derivative, i.e., the exterior differentiation with respect to fiber variables y^i . Using the expressions (2.10), it follows that

$$(2.12) \quad d_V \omega^0 \equiv -\theta^0_a \wedge \omega^a, \quad \text{mod } I\{\omega^0\},$$

where $\theta^0_a = \partial_a \partial_b F dy^b$. As a result of (2.12) one obtains

$$d\omega^0 = -2\theta^0_0 \wedge \omega^0 - \theta^0_a \wedge \omega^a + T_{0i} \omega^0 \wedge \omega^i + \frac{1}{2} T_{ab} \omega^a \wedge \omega^b + T_{an} \omega^a \wedge \omega^n.$$

By setting $\phi_0 := \theta^0_0 - T_{0i} \omega^i$, $\theta_a := \theta^0_a - \frac{1}{2} T_{ab} \omega^b - T_{an} \omega^n$, one arrives at

$$(2.13) \quad d\omega^0 = -2\phi_0 \wedge \omega^0 - \theta_a \wedge \omega^a,$$

which implies that ω^0 has rank $2n-1$, i.e., $\omega^0 \wedge (d\omega^0)^{n-1} \neq 0$.

The 1-forms θ_a define vertical 1-forms with respect to the fibration $\pi : \mathcal{C} \rightarrow M$. The 1-forms θ_a are linearly independent everywhere when the fibers \mathcal{C}_x are tangentially non-degenerate at every point, i.e., $\det(\partial_a \partial_b F) \neq 0$.

Since the vertical 1-forms $\theta_1, \dots, \theta_{n-1}$ are linearly independent, at each point $(x; y) \in \mathcal{C}$, one obtains a coframe $\mathcal{I}_{\text{tot}} := \{\omega^0, \dots, \omega^n, \theta_1, \dots, \theta_{n-1}\}$ for $T_{(x; z)} \mathcal{C}$ adapted to the flag

$$(2.14) \quad 0 \subset VT_p \mathcal{C} \subset K_p \subset H_p \subset T_p \mathcal{C},$$

defined previously in (2.11). Such a coframe is called *1-adapted*. For a quadric, this filtration coincides with the one introduced in (2.8). In analogy with the flat model, the following algebraic ideals are defined:

$$(2.15) \quad \begin{aligned} \mathcal{I}_{\text{tot}} &= I\{\omega^0, \dots, \omega^n, \theta^1, \dots, \theta^{n-1}\}, \\ \mathcal{I}_{\text{bas}} &= I\{\omega^0, \dots, \omega^n\}, \\ \mathcal{I}_{\text{char}} &= I\{\omega^0, \dots, \omega^{n-1}, \theta^1, \dots, \theta^{n-1}\}. \end{aligned}$$

Using the expressions (2.10), equation (2.13) gives

$$(2.16) \quad \theta_a = \partial_a \partial_b F dx^b + \partial_0 \partial_a F dx^0 + \partial_b \partial_a F dx^b + (\partial_a \partial_n F - \partial_a F - \partial_0 F \partial_a F) dx^n$$

From the expression above it follows that the linear independence of θ_a 's modulo ω^i 's is equivalent to the non-degeneracy of the vertical Hessian $\partial_a \partial_b F$.

The fact that such a coframing is adapted to the flag (2.14) and satisfies (2.13) implies that different choices of 1-adapted coframes are related by

$$\begin{aligned} \omega^0 &\mapsto \mathbf{a}^2 \omega^0, & \omega^a &\mapsto \mathbf{E}_a \omega^a + \mathbf{A}^a_b \omega^b, \\ \omega^n &\mapsto \mathbf{e} \omega^0 + \mathbf{B}_a \omega^a + \mathbf{b}^2 \omega^n, & \theta_a &\mapsto \mathbf{C}_a \omega^0 + \mathbf{S}_{ab} (\mathbf{A}^{-1})^b_c \omega^c + \mathbf{a}^2 (\mathbf{A}^{-1})^b_a \theta_b. \end{aligned}$$

In the language of G -structures, the structure group of the bundle of 1-adapted coframes, G_1 , is of the form

$$(2.17) \quad \left\{ \begin{pmatrix} \mathbf{a}^2 & 0 & 0 & 0 \\ \mathbf{E} & \mathbf{A} & 0 & 0 \\ \mathbf{e} & \mathbf{B} & \mathbf{b}^2 & 0 \\ \mathbf{C} & {}^t \mathbf{A}^{-1} \mathbf{S} & 0 & \mathbf{a}^2 {}^t \mathbf{A}^{-1} \end{pmatrix} \mid \begin{array}{ll} \mathbf{A} \in \text{GL}_{n-1}(\mathbb{R}), & \mathbf{a}, \mathbf{b} \in \mathbb{R} \setminus \{0\}, \\ {}^t \mathbf{B}, \mathbf{C}, \mathbf{E} \in \mathbb{R}^{n-1}, & \mathbf{e} \in \mathbb{R}, \\ \mathbf{S} \in M_{n-1}(\mathbb{R}), & \mathbf{S} = {}^t \mathbf{S} \end{array} \right\}$$

Remark 2.5. The reason for using \mathbf{a}^2 and \mathbf{b}^2 in G_1 is as follows. Let $L(x^i; y^i)$ be a Lagrangian function for a causal structure, i.e., the vanishing set of L defines \mathcal{C} . As is conventional in the case of pseudo-conformal structures, it is desirable to restrict to the set of causal transformations preserving the sign of L in a neighborhood of the null cones, i.e., restricting to the set of Lagrangians of the form $S^2 L$ for some nowhere vanishing homogeneous function $S : TM \rightarrow \mathbb{R}$. Recall that the projective Hilbert form in terms of L is given by $\omega^0 = \partial_i L dx^i$. Replacing L by $S^2 L$ and restricting to the zero locus of L , the projective Hilbert form transforms to $S^2(L_i dx^i) = S^2 \omega^0$. This observation justifies the use of \mathbf{a}^2 in G_1 . Moreover, recall that a vector bundle of rank 1 is called orientable if it has a non-vanishing global section. In the case of contact manifolds, a contact distribution is called *co-orientable* or *transversally orientable* if it has a globally defined contact form [LM87]. Similarly, a quasi-contact structure is called co-orientable if it has a globally defined quasi-contact form ω^0 . If the quasi-contact distribution on \mathcal{C} is given a co-orientation, it is preserved under the action of G_1 . Similarly, if the characteristic field of a quasi-contact structure (see Section 2.3.5) is given an orientation, then having \mathbf{b}^2 in (2.17) implies that the structure group G_1 preserves the orientation of the characteristic field. Moreover, the conformal scaling of the quadratic form g defined in 2.3.11 by the action of G_1 is given by $\mathbf{a}^2 \mathbf{b}^2$. As a result, it is possible to define space-like and time-like horizontal vector fields v in $T\mathcal{C}$ via the signature of $g(v, v)$.

To implement Cartan's method of equivalence, one lifts the 1-adapted coframe to the G_1 -bundle in the following way (see [IL03, Gar89, Olv95] for the background). Recall that the set of all coframes on \mathcal{C} , defines a $\text{GL}(2n)$ -principal bundle $\tilde{\zeta} : \tilde{\mathcal{P}} \rightarrow \mathcal{C}$, with the right action

$$R_g(\vartheta_p) = g^{-1} \cdot \vartheta_p,$$

where $\vartheta_p \in \zeta^{-1}(p)$ is a coframe at $p \in \mathcal{C}$, $g \in \text{GL}(2n)$, and the action on the right hand side is the ordinary matrix multiplication.

Among the set of all coframes on \mathcal{C} , let ϑ represent 1-adapted coframes ${}^t(\omega^0, \dots, \theta_{n-1})$, which are well-defined up to the action of G_1 . As a result, the 1-adapted coframes are sections of a G_1 -bundle $\varsigma_1 : \mathcal{P}_1 \rightarrow \mathcal{C}$. On $\mathcal{P}_1 \cong \mathcal{C} \times G_1$ one can define a canonical set of 1-forms by setting

$$\underline{\vartheta}(p, g) = {}^t(\underline{\omega}^0, \dots, \underline{\theta}_{n-1}) := g^{-1} \cdot \vartheta_p.$$

The exterior derivatives of the 1-forms in $\underline{\vartheta}$ are given by

$$(2.18) \quad d\underline{\vartheta} = dg^{-1} \wedge \vartheta + g^{-1} \cdot d\vartheta = -g^{-1} \cdot dg \cdot \wedge g^{-1}\vartheta + g^{-1} \cdot d\vartheta = -\alpha \wedge \underline{\vartheta} + T,$$

where $\alpha(p, g) = g^{-1} \cdot dg$ is \mathfrak{g}_1 -valued 1-form where \mathfrak{g}_1 denotes the Lie algebra of G_1 . The 2-forms $T(p, g)$ can be expressed in terms of $\underline{\omega}^i \wedge \underline{\omega}^j$, $\underline{\theta}_a \wedge \underline{\omega}^i$, $\underline{\theta}_a \wedge \underline{\theta}_b$. The terms constituting T are called the torsion terms. They are semi-basic with respect to the projection ς_1 . Via α , one can identify the vertical tangent spaces of the fibers of $\varsigma_1 : \mathcal{P}_1 \rightarrow \mathcal{C}$ with \mathfrak{g}_1 . The \mathfrak{g}_1 -valued 1-form α is called a *pseudo-connection* on \mathcal{P}_1 which is not yet uniquely defined. Recall that the difference between a *connection* and a pseudo-connection on G_1 -bundle lies in the fact that a connection form $\tilde{\alpha}$ satisfies the equivariance condition $R_g^* \tilde{\alpha} = \text{Ad}(g^{-1}) \tilde{\alpha}$, where $g \in G_1$, while a pseudo-connection in general does not.

A choice of pseudo-connection in (2.18) gives rise to structure equations for the 1-forms $\underline{\vartheta}$. For 1-adapted coframes, the pseudo-connection forms are of the form,

$$(2.19) \quad \begin{pmatrix} 2\phi_0 & 0 & 0 & 0 \\ \gamma^a_0 & \psi^a_b & 0 & 0 \\ \gamma^n_0 & \gamma^n_a & 2\phi_n & 0 \\ \pi_a & \pi_{ab} & 0 & 2\phi_0 \delta^a_b - \psi^a_b \end{pmatrix}.$$

where $\pi_{ab} = \pi_{ba}$.

It follows that

$$(2.20a) \quad d\underline{\omega}^0 = -2\phi_0 \wedge \underline{\omega}^0 - \underline{\theta}_a \wedge \underline{\omega}^a,$$

$$(2.20b) \quad d\underline{\omega}^a = -\gamma^a_0 \wedge \underline{\omega}^0 - \psi^a_b \wedge \underline{\omega}^b - h^{ab} \underline{\theta}_b \wedge \underline{\omega}^n,$$

$$(2.20c) \quad d\underline{\omega}^n = -\gamma^n_0 \wedge \underline{\omega}^0 - \gamma^n_a \wedge \underline{\omega}^a - 2\phi_n \wedge \underline{\omega}^n,$$

$$(2.20d) \quad d\underline{\theta}_a = -\pi_a \wedge \underline{\omega}^0 - \pi_{ab} \wedge \underline{\omega}^b + \psi^b_a \wedge \underline{\theta}_b - 2\phi_0 \wedge \underline{\theta}_a + \sigma_a \wedge \underline{\omega}^n + \sigma^b_a \wedge \underline{\theta}_b,$$

where the 1-forms σ_a, σ^b_a , are the torsion terms and vanish modulo \mathcal{I}_{tot} . Their components are determined in Section 2.3.9.

The equation of $d\omega^0$ follows directly from (2.13). Moreover, the Pfaffian system \mathcal{I}_{bas} , is completely integrable with the fibers \mathcal{C}_x being its integral manifolds. It follows that $d\omega^a = \tau^a_i \wedge \omega^i$, and $d\omega^n = \tau^n_i \wedge \omega^i$, where $\tau^a_i, \tau^n_i \equiv 0$ modulo \mathcal{I}_{tot} . Using the fact that the action of G_1 preserves \mathcal{I}_{bas} , and that two pseudo-connection differ by a semi-basic \mathfrak{g}_1 -valued 1-form, all the 1-forms τ^a_i, τ^n_i can be absorbed into the pseudo-connection forms. The only torsion term left is $h^{ab} \theta_b \wedge \omega^n$, for some function h^{ab} , which is discussed in Section 2.3.6. From now on, since structure equations are always written for the lifted coframe $\underline{\vartheta}$, the underline in $\underline{\omega}^i$ and $\underline{\theta}_a$'s will be dropped.

Remark 2.6. The oriented projectivized tangent bundle $\mathbb{P}_+ TM$ is defined so that at $x \in M$ the fiber $\mathbb{P}_+ T_x M$ is isomorphic to the space of *rays* in $T_x M$, or, equivalently, the set of oriented real lines in $T_x M$ passing through the origin. As a result, $\mathbb{P}_+ T_x M$ is the nontrivial double cover of $\mathbb{P} T_x M$.

The above definition of cone structures, and more specifically causal structures, can be changed so that \mathcal{C} is a sub-bundle of $\mathbb{P}_+ TM$. In the case of causal structures immersed in $\mathbb{P}_+ TM$, the assumption that the fibers $\mathcal{C}_x = \iota^{-1}(x)$ be connected is not desirable. This is due to the fact that \mathcal{C}_x should be thought of as the light cones which are comprised of the future and past light cones whose projectivizations are disjoint. Thus, one expects $\mathcal{C}_x = \mathcal{C}_x^+ \sqcup \mathcal{C}_x^-$. This way, in analogy with Finsler geometry, a cone structure is called *reversible* if $\widehat{\mathcal{C}} = -\widehat{\mathcal{C}}$. Otherwise, it is called *non-reversible*. It should be noted that among oriented causal structures, non-reversible ones can be of interest in certain problems due to possible differences in their future and past light cones. This is similar to the case of non-reversible Finsler structures containing the important class of Randers spaces [BRS04]. In [KP67], Kronheimer and Penrose use the term *self-dual* for a reversible causal structure. This is due to the fact that they define the dual of a causal structure to be the causal structure obtained from replacing all the causal relations between the points of the space by their inverses, i.e., interchanging the words “future” and “past”.

2.3.4. A dual formulation of causal structures. In the context of geometric control theory, geometric structures arising from distributions are typically formulated in the cotangent bundle [AZ02]. Firstly, the definition of a Legendre transformation is recalled.

Definition 2.7. Given a sufficiently smooth real-valued function L , usually referred to as a Lagrangian, on TM , the associated Legendre transformation $\text{Leg} : TM \rightarrow T^*M$, is defined by

$$\text{Leg} = \hat{\pi} \circ d_V L,$$

where $d_V L$ defines a section of the product bundle $TM \times T^*M$, with the projection map $\hat{\pi} : TM \times T^*M \rightarrow T^*M$. In local coordinates, $\text{Leg}(x^i; y^i) = (x^i; p_i)$ where $p_i = \frac{\partial L}{\partial y^i}$, and (p_0, \dots, p_n) are local coordinates for the fibers of T^*M .

The Legendre transformation is a local diffeomorphism provided that $\text{Leg}_* = \frac{\partial^2 L}{\partial y^i \partial y^j}$ is non-degenerate everywhere. The inverse Legendre transformation Leg^{-1} can be obtained by the same procedure using the Hamiltonian function $Q := \mathcal{A} \circ \text{Leg}^{-1}$ where $\mathcal{A} := dL(\underline{A}) - L$ with \underline{A} being the Liouville vector field, i.e., $\mathcal{A} = y^i \frac{\partial L}{\partial y^i} - L$ in local coordinates. When the degree of homogeneity of L is $r \neq 1$, Euler's theorem gives $Q = (r-1)L \circ \text{Leg}^{-1}$.

Changing the defining function to $L'(x, y) = S(x, y)L(x, y)$, the function Leg remains invariant over \mathcal{C} , hence the vanishing set of Q in T^*M remains invariant. Consequently, one can equivalently study the *dual causal structure* $\mathcal{C}^* := \text{Leg}(\mathcal{C}) \subset \mathbb{P}T^*M$ whose fibers $\mathcal{C}_x^* \subset \mathbb{P}T_x^*M$ are the projective dual of the projective hypersurfaces $\mathcal{C}_x \subset \mathbb{P}T_x M$.

Via the Legendre transformation the projective Hilbert form $\omega^0 = \partial_i L dx^i$ is the pull-back of the canonical contact form $\lambda = p_i dx^i$ restricted to $\mathcal{C}^* \subset \mathbb{P}T^*M$ i.e.,

$$\text{Leg}^*(p_i dx^i) = \frac{\partial L}{\partial y^i}(x, y) dx^i.$$

Consequently, the coframe adaptation can be carried out using λ for the dual cone structure. Hence, this dual formulation can be taken as the starting point for studying causal structures.

2.3.5. Characteristic curves and a quasi-Legendrian foliation. A causal structure comes with a distinguished set of curves which can be defined using the quasi-contact nature of \mathcal{C} . Recall from Example 2.2 that a quasi-contact structure (or even contact structure) on a $2n$ -dimensional manifolds was defined locally in terms of a 1-form of rank $2n-1$. As a result, there is a unique direction along which ω^0 and $d\omega^0$ are degenerate i.e., $[\mathbf{v}] \subset T\mathbb{P}TM$ such that $\mathbf{v} \lrcorner \omega^0 = 0$ and $\mathbf{v} \lrcorner d\omega^0 = 0$, which is called the *characteristic field* of the quasi-contact structure and its integral curves are called *characteristic curves* (c.f. [Hsu92]). A section of the characteristic field is called a *characteristic vector field*.

In local coordinates using the defining function F in (2.2), one obtains that the vector field

$$F \frac{\partial}{\partial x^0} + y^a \frac{\partial}{\partial x^a} + \frac{\partial}{\partial x^n} + A^a \frac{\partial}{\partial y^a}$$

where

$$A^b \partial_a \partial_b F = -F \partial_0 \partial_a F - y^b \partial_b \partial_a F + \partial_a F + \partial_0 F \partial_a F - \partial_a \partial_n F$$

represents a characteristic vector field on \mathcal{C} .

Definition 2.8. Let M be a $2n$ -dimensional manifold with a quasi-contact 1-form $\omega \in \Gamma(T^*M)$. A submanifold $N \subset M$ is called *quasi-Legendrian* if $T_x N \in \text{Ker } \omega$ and $T_x N \lrcorner d\omega = 0$ for all $x \in N$, with the property that $T_x N$ is transversal to the characteristic field of ω .

The bundle \mathcal{C} is $2n$ -dimensional and $\omega^0 \in T^*\mathcal{C}$ has maximal rank $2n-1$, therefore, \mathcal{C} has a quasi-contact structure. The fibers \mathcal{C}_x form a quasi-Legendrian foliation of \mathcal{C} since the vertical tangent spaces $T_y \mathcal{C}_x$ are annihilated by $\mathcal{I}_{\text{bas}} = \{\omega^0, \dots, \omega^n\}$.

Moreover, using (2.13), the characteristic field of ω^0 is the kernel of the Pfaffian system $\mathcal{I}_{\text{char}} = \{\omega^0, \dots, \omega^{n-1}, \theta_1, \dots, \theta_{n-1}\}$, which justifies the subscript *char*.

Remark 2.9. Recall that the geodesics of a Finsler structure are defined as the projection of integral curves of the Reeb vector field associated to the contact structure on the indicatrix bundle induced by the Hilbert form [Bry02]. Here, the projection of the characteristic curves gives an analogue of null geodesics in pseudo-conformal structures.

2.3.6. *Projective second fundamental form and the second coframe adaptation.* The next coframe adaptation involves the projective fundamental form of the fibers \mathcal{C}_x .

Let $\omega^a \equiv -h^{ab}\theta_b \wedge \omega^n$ modulo $I\{\omega^0, \dots, \omega^{n-1}\}$. Using the identity $d^2\omega^0 = 0$, it follows that

$$h^{ab}\theta_a \wedge \theta_b \wedge \omega^n = 0,$$

i.e., $h^{ab} = h^{ba}$. Using the local expressions of ω^a in (2.10), one obtains that

$$d_V \omega^a \equiv -dy^a \wedge \omega^n \mod I\{\omega^0, \dots, \omega^{n-1}\}.$$

Recall from (2.12) that $dy^a = h^{ab}\theta_b$ where $(h^{ab}) = (\partial_a \partial_b F)^{-1}$.

Using the structure equations (2.20), the infinitesimal action of the structure group G_1 on h^{ab} is obtained from the identity $d^2\omega^a = 0$, and is given by

$$(2.21) \quad dh^{ab} + h^{ac}\psi^b{}_c + h^{bc}\psi^a{}_c - 2h^{ab}(\phi_0 + \phi_n) \equiv 0 \mod \mathcal{I}_{\text{tot}}.$$

The relation above implies that the functions $h^{ab}(p, g_0)$ and $\tilde{h}^{ab}(p, g_1)$ where $g_1 = gg_0$, with $g \in G_1$ represented by the matrix (2.17), are related by

$$(2.22) \quad \tilde{h}^{ab} = \mathbf{a}^2 \mathbf{b}^2 (\mathbf{A})^{-1a}{}_c (\mathbf{A})^{-1b}{}_d h^{cd}.$$

It follows that the bilinear form $\phi_2 = h^{ab}\theta_a \circ \theta_b$ when pulled-back to the fibers \mathcal{C}_x coincides with its projective second fundamental form.

Remark 2.10. The action of the structure group on h^{ab} can also be obtained directly from the structure equations (2.20). However, in implementing Cartan's method of equivalence finding the infinitesimal group action by differentiating the structure equations is more straightforward and computationally easier. This is why from now on the focus will be primarily on the infinitesimal action of the structure group on the torsion terms.

Note that by definition 2.4, tangential non-degeneracy of the fibers \mathcal{C}_x implies that h^{ab} is non-degenerate everywhere in \mathcal{C} . As a result, the signature of h^{ab} , as a well-defined tensor over \mathcal{C} , remains the same everywhere.

Assuming that h^{ab} has signature (p, q) , the relation (2.22) implies that $\mathbf{A}^a{}_b$ can be chosen so that h^{ab} is normalized to ε^{ab} . By restricting the G_1 -bundle to the coframes for which $h^{ab} = \varepsilon^{ab}$, and using the relation (2.21), it follows that

$$(2.23) \quad \varepsilon^{ac}\phi^b{}_c + \varepsilon^{bc}\phi^a{}_c \equiv 0 \mod \mathcal{I}_{\text{tot}}, \quad \text{where} \quad \phi^a{}_b = \psi^a{}_b - (\phi_0 + \phi_n)\delta^a{}_b.$$

As a result, for such coframes one can replace $\psi^a{}_b$ by $\phi^a{}_b + (\phi_0 + \phi_n)\delta^a{}_b$ where the matrix-valued 1-forms $\phi^a{}_b$ take values in $\mathfrak{o}(p, q)$ modulo \mathcal{I}_{tot}

Coframings with normalized h^{ab} are *2-adapted*, and they are local sections of a G_2 -bundle over \mathcal{C} where

$$(2.24) \quad G_2 = \left\{ \begin{pmatrix} \mathbf{a}^2 & 0 & 0 & 0 \\ \mathbf{E} & \mathbf{a}\mathbf{b}\mathbf{A} & 0 & 0 \\ \mathbf{e} & \mathbf{B} & \mathbf{b}^2 & 0 \\ \mathbf{C} & \mathbf{A}\mathbf{S} & 0 & \frac{\mathbf{a}}{\mathbf{b}}\mathbf{A} \end{pmatrix} \left| \begin{array}{ll} \mathbf{A} \in \text{O}(p, q), & \mathbf{a}, \mathbf{b} \in \mathbb{R} \setminus \{0\}, \\ {}^t\mathbf{B}, \mathbf{C}, \mathbf{E} \in \mathbb{R}^{n-1}, & \mathbf{e} \in \mathbb{R}, \\ \mathbf{S} \in M_{n-1}(\mathbb{R}), & \mathbf{S} = {}^t\mathbf{S} \end{array} \right. \right\}.$$

Remark 2.11. By a result of Griffiths and Harris [GH79], if the projective second fundamental form of a projective hypersurface $\mathcal{C}_x \subset \mathbb{P}T_x M$ has degeneracy of rank k , then its projective dual $\mathcal{C}_x^* \subset \mathbb{P}T_x^* M$ has codimension $k+1$. In other words, let (M, \mathcal{C}) be a cone structure (M, \mathcal{C}) of codimension one, whose fibers have projective second fundamental form with degeneracy of rank k . Then, its dual cone structure (M, \mathcal{C}^*) has fibers of dimension $n-k-1$. Therefore, it is more natural to study cone structure of codimension one whose fibers have degenerate second fundamental form via their dual formulation since $\dim \mathcal{C} = 2n$, and $\dim \mathcal{C}^* = 2n-k$.

2.3.7. *The third coframe adaptation.* The 1-forms $\phi^a{}_b$ are $\mathfrak{o}(p, q)$ -valued, i.e., $\phi_{ab} + \phi_{ba} \equiv 0$ modulo \mathcal{I}_{tot} , where $\phi_{ab} = \varepsilon_{ac}\phi^c{}_b$. From (2.23) it follows that

$$(2.25) \quad \psi_{ab} = \phi_{ba} + (\phi_0 + \phi_n)\delta_{ab} + E_{aib}\omega^i + F_a{}^c{}_b\theta_c,$$

where $E_{aib} = E_{bia}$ and $F_a{}^c{}_b = F_b{}^c{}_a$. Hence, with respect to 2-adapted coframes, the structure equations are

$$(2.26a) \quad d\omega^0 = -2\phi_0 \wedge \omega^0 - \theta_a \wedge \omega^a,$$

$$(2.26b) \quad d\omega^a = -\gamma^a{}_0 \wedge \omega^0 - \phi^a{}_b \wedge \omega^b - (\phi_0 + \phi_n) \wedge \omega^a - \varepsilon^{ab} \theta_b \wedge \omega^n, \\ - E^a{}_{ic} \omega^i \wedge \omega^c - F^{ab}{}_c \theta_b \wedge \omega^c,$$

$$(2.26c) \quad d\omega^n = -\gamma^n{}_0 \wedge \omega^0 - \gamma^n{}_a \wedge \omega^a - 2\phi_n \wedge \omega^n,$$

$$(2.26d) \quad d\theta_a = -\pi_a \wedge \omega^0 - \pi_{ab} \wedge \omega^b + \phi^b{}_a \wedge \theta_b - (\phi_0 - \phi_n) \wedge \theta_a \\ + \sigma_a \wedge \omega^n + \sigma^b{}_a \wedge \theta_b,$$

where ε^{ab} is used to raise the indices in E_{aib} and $F_a{}^c{}_b$. By replacing

$$\phi^a{}_b \mapsto \phi^a{}_b + \frac{1}{2} (E^a{}_{bc} \omega^c - \varepsilon^{ae} E^d{}_{ec} \varepsilon_{da} \omega^c + \varepsilon_{cd} E^c{}_{ab} \omega^d) \\ \phi_n \mapsto \phi_n + E^b{}_{nb} \omega^n, \quad \gamma^a{}_0 \mapsto \gamma^a{}_0 + E^a{}_{0b} \omega^b,$$

it can be assumed that $E^a{}_{bc}$ and $E^a{}_{0c}$ vanish, and $E^a{}_{nb}$ is trace-free.

Differentiating $d\omega^a$, and collecting the terms of the form $\tau^a{}_b \wedge \omega^b \wedge \omega^n$ for some 1-form satisfying $\tau_{ab} = \tau_{ba}$ and $\tau^a{}_a \equiv 0$ modulo \mathcal{I}_{tot} , one obtains

$$[(dE_{anb} + 2E_{anb}\phi_n + 2E_{cn(a}\phi^c{}_b) + \pi_{ab}) \wedge \omega^n - d(\phi_0 + \phi_n)\varepsilon_{ab}] \wedge \omega^b = 0,$$

where $E_{anb} = \varepsilon_{ac} E^c{}_{nb}$. Contracting the above equations with ε^{ab} , and using the properties $E_{anb} = E_{bna}$, $E^a{}_{na} = 0$, and $\phi_{ab} = -\phi_{ba}$, it follows that $d(\phi_0 + \phi_n) \equiv \frac{1}{n-1} \pi_{ab} \varepsilon^{ab} \wedge \omega^n$ modulo $\mathcal{I}_{\text{char}}$. As a result, the infinitesimal action of the structure group on $E^a{}_{nb}$ is given by

$$(2.27) \quad dE_{anb} + 2E_{anb}\phi_n + 2E_{cn(a}\phi^c{}_b) + (\pi_{ab} - \frac{1}{n-1} \pi_{cd} \varepsilon^{cd} \varepsilon_{ab}) \equiv 0, \quad \text{mod } \mathcal{I}_{\text{tot}}.$$

Similarly to Section 2.3.6, it is possible to restrict to coframes with vanishing $E^a{}_{nb}$, and consequently, by relation (2.27), the 1-forms π_{ab} are pure trace modulo \mathcal{I}_{tot} , i.e., $\pi_{ab} = \pi_n \varepsilon_{ab}$, modulo \mathcal{I}_{tot} , for some 1-form π_n . Such coframes are called *3-adapted*.

The 3-adapted coframes are local sections of a G_3 -bundle, where G_3 consists of matrices

$$(2.28) \quad \left\{ \begin{pmatrix} \mathbf{a}^2 & 0 & 0 & 0 \\ \mathbf{E} & \mathbf{a}\mathbf{b}\mathbf{A} & 0 & 0 \\ \mathbf{e} & \mathbf{B} & \mathbf{b}^2 & 0 \\ \mathbf{C} & \mathbf{c}^t\mathbf{A}\varepsilon & 0 & \frac{\mathbf{a}}{\mathbf{b}}\mathbf{A} \end{pmatrix} \left| \begin{array}{l} \mathbf{A} \in \text{O}(p, q), \quad \mathbf{a}, \mathbf{b} \in \mathbb{R} \setminus \{0\}, \\ {}^t\mathbf{B}, \mathbf{C}, \mathbf{E} \in \mathbb{R}^{n-1}, \quad \mathbf{c}, \mathbf{e} \in \mathbb{R} \end{array} \right. \right\}.$$

2.3.8. Symmetries of F^{abc} . In order to show F^{abc} is completely symmetric, one first differentiates equation (2.26b) to obtain

$$0 \equiv \varepsilon^{ab} (d\theta_b - \phi^c{}_b \wedge \theta_c + (\phi_0 - \phi_n) \theta_b + F^{cd}{}_b \theta_d \wedge \theta_c) \wedge \omega^n, \quad \text{mod } \omega^0, \dots, \omega^{n-1}.$$

Differentiating equation (2.26a) yields,

$$0 = (d\theta_c - \phi^a{}_c \wedge \theta_a + (\phi_0 - \phi_n) \theta_c + F^{ab}{}_c \theta_a \wedge \theta_b) \wedge \omega^c, \quad \text{mod } I\{\omega^0, \omega^n\}.$$

Comparing the two equations above, one obtains, $F^{ab}{}_c \theta_a \wedge \theta_b = 0$, i.e., $F^{ab}{}_c = F^{ba}{}_c$. Recall, in (2.25) the quantities $F^{ab}{}_c$ satisfy $F^{abc} = F^{cba}$, where $F^{abc} = \varepsilon^{cd} F^{ab}{}_d$. As a result, F^{abc} is totally symmetric.

2.3.9. The fourth coframe adaptation, Fubini cubic form and the WSF curvature. The next coframe adaptation reveals the essential invariants of a causal structure, which are the Fubini cubic forms of the fibers \mathcal{C}_x and the Weyl shadow flag curvature (also referred to as the WSF curvature).

Differentiating (2.26b) and collecting terms of the form $\tau^{ab}{}_c \wedge \theta_b \wedge \omega^c$ satisfying $\tau_a{}^b{}_c = \tau_c{}^b{}_a$, one has

$$(2.29) \quad 0 \equiv [(dF^{abc} + F^{abd} \phi^c{}_d + F^{adc} \phi^b{}_d + F^{dbc} \phi^a{}_d - F^{abc}(\phi_0 - \phi_n) \\ - (\gamma^n{}_d \varepsilon^{dc} \varepsilon^{ab} - \gamma^a{}_0 \varepsilon^{bc})) \wedge \theta_b - d(\phi_0 + \phi_n) \varepsilon^{ac}] \wedge \omega_c \quad \text{mod } \omega^0, \omega^n$$

Contacting by ε_{ab} , and defining

$$F^a := \varepsilon_{bc} F^{abc},$$

one arrives at

$$d(\phi_0 + \phi_n) = \frac{1}{n-1} (\gamma^b{}_0 - \gamma^n{}_d \varepsilon^{db} + dF^b) \wedge \theta_b.$$

Putting this back in (2.29), and contracting with ε_{bc} gives

$$dF^a \equiv -F^b \phi^a_b + F^a(\phi_0 - \phi_n) - \frac{n(n-2)}{n-1}(\gamma^a_0 - \gamma^n_b \varepsilon^{ba}) \mod \mathcal{I}_{\text{tot}}.$$

As before, the infinitesimal action above makes it possible to translate F^a to zero. Therefore, the class of 4-adapted coframes can be defined by the *apolarity relation*

$$(2.30) \quad F^a = 0.$$

The resulting reduction in the structure group is realized via the relation $\gamma^a_0 \equiv \varepsilon^{ab} \gamma^n_b$ modulo \mathcal{I}_{tot} .

It can be shown that the cubic form

$$F_3 = F^{abc} \theta_a \circ \theta_b \circ \theta_c,$$

when pulled-back to the fiber \mathcal{C}_x , coincides with Fubini cubic form of \mathcal{C}_x . This becomes evident in the derivation of the local expression of F^{abc} in (2.65).

It is noted that because the second fundamental form is normalized, for any choice of 4-adapted coframe, the Fubini cubic form is well-defined tensor on \mathcal{C} .

The structure group G_4 is reduced to the group of matrices of the form

$$(2.31) \quad \left\{ \begin{pmatrix} \mathbf{a}^2 & 0 & 0 & 0 \\ \mathbf{E} & \mathbf{a}\mathbf{b}\mathbf{A} & 0 & 0 \\ \mathbf{e} & \mathbf{B} & \mathbf{b}^2 & 0 \\ \mathbf{C} & \mathbf{c}^t\mathbf{A}\varepsilon & 0 & \frac{\mathbf{a}}{\mathbf{b}}^t\mathbf{A} \end{pmatrix} \left| \begin{array}{l} \mathbf{A} \in \text{O}(p, q), \quad \mathbf{a}, \mathbf{b} \in \mathbb{R} \setminus \{0\}, \\ {}^t\mathbf{B}, \mathbf{C} \in \mathbb{R}^{n-1}, \quad \mathbf{c}, \mathbf{e} \in \mathbb{R} \\ \mathbf{E} = \frac{\mathbf{a}}{\mathbf{b}} \mathbf{A} {}^t\mathbf{B} \end{array} \right. \right\},$$

where ${}^t\mathbf{B}$ denotes the transpose of \mathbf{B} with respect to ε .

By setting $\gamma^a_0 - \varepsilon^{ab} \gamma^n_b = K^a_{i0} \omega^i + K^{ab} \theta_b$, the structure equations after the fourth coframe adaptation are of the form

$$(2.32a) \quad d\omega^0 = -2\phi_0 \wedge \omega^0 - \theta_a \wedge \omega^a,$$

$$(2.32b) \quad d\omega^a = -\varepsilon^{ab} \gamma_b \wedge \omega^0 - \phi^a_b \wedge \omega^b - (\phi_0 + \phi_n) \wedge \omega^a - \varepsilon^{ab} \theta_b \wedge \omega^n, \\ - K^a_{i0} \omega^i \wedge \omega^0 - K^{ab} \theta_b \wedge \omega^0 - F^{ab}_c \theta_b \wedge \omega^c,$$

$$(2.32c) \quad d\omega^n = -\gamma^n_0 \wedge \omega^0 - \gamma_a \wedge \omega^a - 2\phi_n \wedge \omega^n,$$

$$(2.32d) \quad d\theta_a = -\pi_a \wedge \omega^0 - \varepsilon_{ab} \pi_n \wedge \omega^b + \phi^b_a \wedge \theta_b - (\phi_0 - \phi_n) \wedge \theta_a \\ + A^b_a \theta_b \wedge \omega^n + W_{anbn} \omega^b \wedge \omega^n + B^{bc}_a \theta_b \wedge \theta_c \\ + E_{an}{}^c{}_b \theta_c \wedge \omega^b + W_{anbc} \omega^b \wedge \omega^c$$

where γ^n_a is being replaced by γ_a .

Differentiating (2.32a) yields

$$A^a_b = 0, \quad W_{anbn} = W_{bnan}, \quad E_{anbc} = E_{ancb}, \\ K^{ab} = K^{ba}, \quad B^{ab}_c = 0, \quad W_{[a|n|bc]} = 0$$

By making the following changes

$$\gamma_a \mapsto \gamma_a - K_{cn0} \omega^n - K_{(ab)0} \omega^a, \quad \phi_n \mapsto \phi_n - K_{cn0} \omega^c, \\ \phi^a_b \mapsto \phi^a_b + \frac{1}{2} (T^a_{bc} \omega^c - \varepsilon^{ae} T^d_{ec} \varepsilon_{da} \omega^c + \varepsilon_{cd} T^c_{ab} \omega^d) + \frac{1}{2} (K^a_b - \varepsilon^{ac} K^d_c \varepsilon_{eb}) \omega^0,$$

where $T^a_{bc} = \delta^a_b K_{cn0} - \delta^a_c K_{bn0}$, and $K_{an0} = \varepsilon_{ab} K^b_{n0}$, it can be assumed K^a_{i0} vanishes

By replacing

$$\pi_n \mapsto \pi_n + \frac{1}{n-1} \varepsilon^{ac} E_{anbc} \theta^b - \frac{1}{n-1} \varepsilon^{cd} W_{cndn} \omega^n,$$

the traces of W_{anbn} and E^a_{nbc} are absorbed in π_n , and therefore, W_{anbn} and E_{anbc} can be chosen to be trace-free in (a, b) .

The symmetric trace-free tensor W_{anbn} is called the *Weyl shadow flag curvature* (the WSF curvature), which, as will be shown, is the second essential invariant of causal structures.

By differentiating (2.32b), and collecting 3-forms $A_{ab} \theta^a \wedge \omega^b \wedge \omega^n$ such that $A_{ab} = A_{ba}$, and $A^a_a = 0$, it follows that

$$E_{anbc} = -f_{abc},$$

where $f_{abc} = F_{abc;n}$, i.e., the derivative of F_{abc} along the characteristic curves.

2.3.10. *The fifth coframe adaptation.* There is one more coframe adaptation that can be imposed on any causal structure, which reduces the structure group by one dimension. First, note that the derivative of (2.32b) gives

$$(2.33) \quad \left((dK^{ab} + 2K^{c(a}\phi^{b)})_c - 2K^{ab}(\phi_0 - \phi_n) - \varepsilon^{ab}\gamma^n{}_0 - F^{abc}\gamma_c \right) \wedge \theta_b + d\gamma^a \wedge \omega^0 \equiv 0$$

modulo $I\{\omega^1, \dots, \omega^n\}$. To find the infinitesimal group action on K^{ab} via the relation above one needs to find the 2-form components $\nu^{ab} \wedge \theta_b$ of $d\gamma^a$ where $\nu^{ab} \neq 0$ modulo \mathcal{I}_{tot} . Differentiating (2.32c), it follows that

$$(d\gamma_a + F^{bc}{}_a \gamma_b \wedge \theta_c + \Lambda_a) \wedge \omega^a \equiv 0$$

where Λ_a is not of the form $\nu^b{}_a \wedge \theta_b$ with $\nu^b{}_a \neq 0$ modulo \mathcal{I}_{tot} . Inserting the above expression of $d\gamma_a$ into (2.33), one sees that the infinitesimal action of the structure group G_5 on K^{ab} is given by

$$dK^{ab} + 2K^{c(a}\phi^{b)}{}_c - 2K^{ab}(\phi_0 + \phi_n) + \varepsilon^{ab}\gamma^n{}_0 - 2F^{abc}\gamma_c \equiv 0 \quad \text{mod } \mathcal{I}_{\text{tot}}$$

Note that K^{ab} is not a tensor as its infinitesimal transformation rule in terms of the structure group involves $F^{ab}{}_c$.

Using the apolarity relation (2.30), and the fact that K_{ab} and ϕ_{ab} are symmetric and skew-symmetric respectively, the infinitesimal action of the structure group G_4 on $K := \varepsilon_{ab}K^{ab}$ is found to be

$$dK + K(\phi_0 - \phi_n) + (n-1)\gamma^n{}_0 \equiv 0, \quad \text{mod } \mathcal{I}_{\text{tot}}.$$

Now, the 5-adapted coframes can be defined by the property that $K = 0$. Hence, one obtains

$$\gamma^0{}_n = M_{ni}\omega^i + L^a\theta_a.$$

By replacing

$$\gamma_a \mapsto \gamma_a - M_{na}\omega^0 - \varepsilon_{ab}M_{nn}\omega^b, \quad \phi_n \mapsto \phi_n - \frac{1}{2}M_{nn}\omega^0,$$

it can be assumed that $M_{ni} = 0$.

The 5-adapted coframes are local sections of a G_5 -bundle where G_5 is the matrix group defined by

$$(2.34) \quad \left\{ \begin{pmatrix} \mathbf{a}^2 & 0 & 0 & 0 \\ \mathbf{E} & \mathbf{a}\mathbf{b}\mathbf{A} & 0 & 0 \\ \mathbf{e} & \mathbf{B} & \mathbf{b}^2 & 0 \\ \mathbf{C} & \mathbf{c}^t\mathbf{A}\varepsilon & 0 & \frac{\mathbf{a}}{\mathbf{b}}{}^t\mathbf{A} \end{pmatrix} \left| \begin{array}{l} \mathbf{a}, \mathbf{b} \in \mathbb{R} \setminus \{0\}, \mathbf{c} \in \mathbb{R}, \\ \mathbf{A} \in \text{O}(p, q), \mathbf{B}, \mathbf{C} \in \mathbb{R}^{n-1} \\ \mathbf{E} = \frac{\mathbf{a}}{\mathbf{b}}\mathbf{A}^t\mathbf{B}, \mathbf{e} = \frac{1}{2\mathbf{b}^2}\mathbf{B}^t\mathbf{B} \end{array} \right. \right\}.$$

2.3.11. *Invariant conformal quadratic form.* According to (2.34), the structure group G_5 preserves the conformal class of the quadratic form

$$(2.35) \quad g := 2\omega^0 \circ \omega^n - \varepsilon_{ab}\omega^a \circ \omega^b,$$

which is semi-basic with respect to the fibration $\pi : \mathcal{C} \rightarrow M$. As will be shown in Section 2.3.14, the vanishing of the Fubini form implies that the null cones of the causal structure coincide with those of g . The $\mathfrak{o}(p, q)$ -valued pseudo-connection associated to the structure group G_5 acting on g , is given by

$$\begin{pmatrix} \phi_n - \phi_0 & \theta & 0 \\ {}^t\gamma & \phi & {}^t\theta \\ 0 & \gamma & \phi_0 - \phi_n \end{pmatrix},$$

where ${}^t\gamma$ and ${}^t\theta$ are column vectors of 1-forms $\gamma^a := \varepsilon^{ab}\gamma_b$ and $\theta^a := \varepsilon^{ab}\theta_b$, respectively. From now on, ε_{ab} will be used to raise and lower the indices a, b, c .

2.3.12. *The first order structure equations.* Having carried out five coframe adaptations, the first structure equations for causal structures can now be expressed as follows.

The pseudo-connection form for the first order structure group is

$$(2.36) \quad \begin{pmatrix} 2\phi_0 & 0 & 0 & 0 \\ \gamma^a & \phi^a{}_b + (\phi_0 + \phi_n)\delta^a{}_b & 0 & 0 \\ 0 & \gamma_b & 2\phi_n & 0 \\ \pi_a & \varepsilon_{ab}\pi_n & 0 & -\phi^a{}_b + (\phi_0 - \phi_n)\delta^a{}_b \end{pmatrix}.$$

Consequently, the structure equations can be written as

$$\begin{aligned}
(2.37a) \quad d\omega^0 &= -2\phi_0 \wedge \omega^0 - \theta_a \wedge \omega^a, \\
(2.37b) \quad d\omega^a &= -\gamma^a \wedge \omega^0 - \phi^a_b \wedge \omega^b - (\phi_0 + \phi_n) \wedge \omega^a - \theta^a \wedge \omega^n \\
&\quad - K^a_b \theta^b \wedge \omega^0 - F^a_{bc} \theta^b \wedge \omega^c, \\
(2.37c) \quad d\omega^n &= -\gamma_a \wedge \omega^a - 2\phi_n \wedge \omega^n - L^a \theta_a \wedge \omega^0 \\
(2.37d) \quad d\theta_a &= -\pi_a \wedge \omega^0 - \pi_n \wedge \omega_a + \phi^b_a \wedge \theta_b - (\phi_0 - \phi_n) \wedge \theta_a \\
&\quad - f_{abc} \theta^b \wedge \omega^c + W_{anbn} \omega^b \wedge \omega^n + \frac{1}{2} W_{anbc} \omega^b \wedge \omega^c,
\end{aligned}$$

with the following symmetries

$$\begin{aligned}
(2.38) \quad F_{abc} &= F_{(abc)}, \quad F^a_{ab} = 0, \quad K^{ab} = K^{ba}, \quad K^a_a = 0, \\
W_{anbn} &= W_{bnan}, \quad W^a_{nan} = 0, \quad W_{anbc} = -W_{0acb}, \quad W_{[a|n|bc]} = 0.
\end{aligned}$$

The infinitesimal group action on the torsion coefficients is equivalent to saying that the following 1-forms are semi-basic with respect to the fibration $\pi : \mathcal{C} \rightarrow M$.

$$\begin{aligned}
(2.39a) \quad DF_{abc} &:= dF_{abc} - F_{abd} \phi^d_c - F_{adc} \phi^d_b - F_{dbc} \phi^d_a - F_{abc} (\phi_0 - \phi_n), \\
(2.39b) \quad DK_{ab} &:= dK_{ab} - K_{ac} \phi^c_b - K_{bc} \phi^c_a - 2K_{ab} (\phi_0 - \phi_n) - 2F_{abc} \gamma^c, \\
(2.39c) \quad DL_a &:= dL_a - L_b \phi^b_a - 3L^a (\phi_0 - \phi_n) + K^{ab} \gamma_b, \\
(2.39d) \quad DW_{anbn} &:= dW_{anbn} - W_{ancn} \phi^c_b - W_{bncn} \phi^c_a - 4W_{anbn} \phi_n, \\
(2.39e) \quad DW_{anbc} &:= dW_{anbc} - W_{dnbc} \phi^d_a - W_{andc} \phi^d_b - W_{anbd} \phi^d_c, \\
&\quad - W_{anbc} (\phi_0 - 3\phi_n) + (W_{anbn} \varepsilon_{cd} + W_{ancn} \varepsilon_{bd}) \gamma^d.
\end{aligned}$$

By the infinitesimal actions (2.39), it follows that no more reduction of the structure group is possible, unless certain non-vanishing conditions on the torsion coefficients are assumed. Such special cases are considered in subsequent sections.

2.3.13. *An $\{e\}$ -structure.* Using the first order structure equations, an $\{e\}$ -structure can be associated to causal structures. The necessary computations are carried out in the Appendix.

Theorem 2.12. *To a causal structure $(M^{n+1}, \mathcal{C}^{2n})$, $n \geq 4$, one can associate an $\{e\}$ -structure on the principal bundle $P_{1,2} \rightarrow \mathcal{P} \rightarrow \mathcal{C}$ of dimension $\frac{(n+3)(n+2)}{2}$ where $P_{1,2} \subset O(p+2, q+2)$ is the parabolic subgroup defined in (2.5). If $n = 3$, then $P_{1,2}$ is replaced by $P_{1,2,3}$. The essential invariants of a causal structure are the Fubini cubic form of the fibers and the WSF curvature. The vanishing of the essential invariants implies that the $\{e\}$ -structure coincides with the Maurer-Cartan forms of $(p+2, q+2)$. The structure equations can be written as*

$$(2.40) \quad d\phi + \phi \wedge \phi = \begin{pmatrix} -\Phi_0 - \Phi_n & -\Pi_n & -\Pi_b & -\Pi_0 & 0 \\ \Omega^n & \Phi_n - \Phi_0 & \Gamma_b & 0 & \Pi_0 \\ \Omega^a & \Theta^a & \Phi^a_b & \Gamma^a & -\Pi^a \\ \Omega^0 & 0 & \Theta_b & \Phi_0 - \Phi_n & \Pi_n \\ 0 & -\Omega^0 & \Omega_b & -\Omega^n & +\Phi_0 + \Phi_n \end{pmatrix},$$

where ϕ is defined in (2.7) and the right hand side of (2.40) is expressed in (A.17).

The proof of the above theorem will be continued in the Appendix and involves lots of tedious calculations. Of course, it is desirable to find the right change of basis so that the $\{e\}$ -structure defines a Cartan connection. This issue is discussed in Section 2.4.3. It is noted the number of torsion elements on the right hand side of the first order structure equations is minimal due to the special absorption process that was undertaken here. When obtaining the $\{e\}$ -structure in the Appendix, to simplify computations, one term will be added to Equation (2.37d) (see Equation A.1.4).

An immediate corollary of Theorem 2.12 is the following.

Corollary 2.13. *The maximal dimension of the symmetry algebra of causal structures on a $(n+1)$ -dimensional manifolds is $\frac{(n+3)(n+2)}{2}$ which occurs only in the flat model.*

Proof. As explained in Section 2.2 the flat model has the maximal symmetries. Now, suppose there is non-flat model that is maximally symmetric. Then, either the Fubini form or the WSF have a non-zero entry. However, by the infinitesimal group actions (2.39), the structure group acts by scaling on both of the essential invariants. As a result, the symmetry group of the causal structure has to be a proper sub-group of $O(p+1, q+1)$ obtained after normalizing the value of that non-zero entry of the essential invariants. Thus, it cannot be maximally symmetric. \square

The corollary above is clear from the parabolic point of view, since any parabolic geometry (G, P) has maximal symmetry dimension equal to $\dim G$ which can only be reached in the flat case [ČS09].

2.3.14. *The case of conformal structures.* Using the first order structure equations, an observation is made regarding (pseudo-)conformal structures viewed as causal structures.

Proposition 2.14. *A causal structure (M, \mathcal{C}) of signature $(p+1, q+1)$ with vanishing Fubini cubic form induces a pseudo-conformal structure of the same signature on M . The resulting pseudo-conformal structure is flat if the initial causal structure has vanishing WSF curvature, and hence flat. Conversely, any pseudo-conformal structure defines a causal structure of the same signature on its bundle of null cones. The resulting causal structure is flat if the initial pseudo-conformal structure is flat.*

Proof. To show the first part, note that if the Fubini form vanishes then, by the Bianchi identities (A.19), the quantities K_{ab} and L_a vanish as well. The first order structure equations (2.36) can be used to show that the Lie derivatives of the quadratic form g , defined in Section 2.3.11, along vertical vector fields $\frac{\partial}{\partial \theta_1}, \dots, \frac{\partial}{\partial \theta_{n-1}}$ are given by

$$(2.41) \quad \mathcal{L}_{\frac{\partial}{\partial \theta_a}} g = \lambda g,$$

for some non-vanishing function λ on \mathcal{C} . The vector fields $\frac{\partial}{\partial \omega^i}$, and $\frac{\partial}{\partial \theta_a}$ are defined as dual to the coframe $\omega^0, \dots, \theta_{n-1}$. The vertical vector fields are integrable with the fibers \mathcal{C}_x as their integral manifolds, and M as the quotient space. Consequently, the conformal class of the quadratic form g descends to M and because it is non-degenerate and of maximal rank on M , a pseudo-conformal structure on M is obtained with the same signature as that of g . If the causal structure is flat, i.e., the $\{e\}$ -structure is given by the Maurer-Cartan forms of $O(p+2, q+2)$, then the structure bundle fibers over M as well, implying that the resulting pseudo-conformal structure has to be flat too.

The converse part is shown similarly. Given a pseudo-conformal structure of signature $(p+1, q+1)$, choose a coframe $(\omega^0, \dots, \omega^n)$ so that a representative of the pseudo-conformal structure takes the form $g = 2\omega^0 \circ \omega^n - \varepsilon_{ab}\omega^a \circ \omega^b$. On its null cone bundle $\pi: \mathcal{C} \rightarrow M$, the metric g pulls back to define a quadratic form on \mathcal{C} satisfying (2.41). The 1-forms ω^i define a set of semi-basic 1-forms on \mathcal{C} . Using conformal transformations, ω^i 's can be chosen to become adapted to the flag (2.11). This can be seen as follows. The affine tangent space $\widehat{T}_y \mathcal{C}_x \subset T_x M$ is a null hyperplane and by a conformal transformation ω^0 can be taken to be its annihilator. Moreover, the dual of ω^0 with respect to the metric g lies on \hat{y} . More explicitly, let $y \in \hat{y} \subset \widehat{\mathcal{C}}$. Because, $g_{ij}y^i y^j = 0$, the 1-form ω^0 as the annihilator of $\widehat{T}_y \mathcal{C}_x$ can be expressed as $\omega^0 = g_{ij}y^j dx^i$. Now, it is apparent that the dual of ω^0 lies in \hat{y} . Since the metric is expressed as $g = 2\omega^0 \circ \omega^n - \varepsilon_{ab}\omega^a \circ \omega^b$, it follows that $\omega^a(\hat{y}) = 0$, and $\omega^n(\hat{y}) \neq 0$. Hence, ω^i 's are adapted to the flag (2.11).

Since g satisfies (2.41), the resulting causal structure have vanishing Fubini form. If the conformal structure is flat, i.e., isomorphic to the flat model, then as discussed in Section 2.2, the corresponding causal structure is flat as well. Because the construction is reversible, the result follows from the first part of the proposition. \square

The following corollary follows immediately.

Corollary 2.15. *If a pseudo-conformal structure lifts to a causal structure with vanishing WSF curvature, then the resulting causal structure is flat.*

Proof. By Proposition 2.14, a pseudo-conformal structure defines a causal structure with vanishing Fubini cubic form. If the resulting causal structure has vanishing WSF curvature, then all of its essential invariants are zero and is therefore flat. By Proposition 2.14, one obtains that the initial pseudo-conformal structure has to be flat as well. \square

Remark 2.16. As was mentioned previously, the discussion above goes through in the case of conformal structures of positive definite signature. The extra step would be to complexify the inner product and the tangent bundle and work with complex causal structures endowed with a reality condition. See [LeB83] for a treatment of complex conformal structures.

2.4. Causal structures as parabolic geometries. This section provides the necessary definitions and theorems needed to characterize causal structures as parabolic geometries. A main reference on parabolic geometries is [ČS09], but, for the purposes of this section the articles [KT14, ANN15, Yam93, Zel09] suffice.

The main step is the characterization of causal structures by a bracket generating distribution with certain properties, described in Section 2.4.2. As a result, causal structures on an $(n + 1)$ -dimensional manifold can be viewed as regular normal infinitesimal flag structures of type $(B_n, P_{1,2})$ or $(D_n, P_{1,2})$ if $n \geq 4$, or $(D_3, P_{1,2,3})$ if $n = 3$. By a result of Tanaka-Morimoto-Čap-Schichl, such structures can be canonically identified as regular normal parabolic geometries of the same type.

2.4.1. Parabolic geometries and Tanaka Theory. A parabolic geometry is a Cartan geometry (\mathcal{P}, ω) of type (G, P) where G is a real or complex semi-simple Lie group and $P \subset G$ is a parabolic subgroup. Recall that a Cartan geometry (\mathcal{P}, ω) of type (G, H) is a principle bundle $H \rightarrow \mathcal{P} \rightarrow M$, endowed with a Cartan connection ω . The 1-form ω takes value in \mathfrak{g} , the Lie algebra of G , and satisfies the following conditions.

- (1) $\omega_u : T_u \mathcal{P} \rightarrow \mathfrak{g}$ is an isomorphism of vector space for all $u \in \mathcal{P}$.
- (2) $V^\dagger \lrcorner \omega = V$ for all $V \in \mathfrak{h}$ where V^\dagger is the fundamental field of V .
- (3) $(R_h)^* \omega = \text{Ad}(h^{-1})\omega$ for all $h \in H$.

A canonical way of identifying a geometric structure as a parabolic geometry is by characterizing it in terms of certain *bracket generating distributions* possibly equipped with a tensor field, e.g., conformal structures or semi-Riemannian structures. Recall the following definitions. To any distribution $D \subset TN$, one can associate a Pfaffian system

$$D^\perp := \{\omega \in T^* \mathcal{C} \mid \omega(u) = 0 \ \forall \ u \in D\}.$$

Given a distribution $D \subset TC$, its space of sections is denoted by $\Gamma(D)$ and its *derived distribution* is defined by $D^2 = D + [D, D]$. Here, $D + [D, D]$ means a distribution whose sections are obtained from $\Gamma(D) + [\Gamma(D), \Gamma(D)]$. Define $D^1 = D$. The k -th *weak derived distribution* of D is defined by

$$(2.42) \quad D^{k+1} = D^k + [D^k, D].$$

One obtains that

$$(2.43) \quad (D^{k+1})^\perp = \{\omega \in (D^k)^\perp \mid d\omega \equiv 0 \ \text{mod} \ (D^k)^\perp\}.$$

A distribution $D \subset TN$ is called *bracket generating* if there exists $k > 1$ such that $D^k = TN$. The small growth vector of a bracket generating distribution D is (d_1, \dots, d_k) where $d_k = \dim(D^k)$.

On a manifold M , a bracket generating distribution, D , of depth μ induces a filtration of $T_x M$ for all $x \in M$, by

$$D(x) := D^1(x) \subset D^2(x) \subset \dots \subset D^\mu(x) = T_x M.$$

Let $\mathfrak{m}(x)$ denote the grading of $T_x M$ that corresponds to the filtration above, i.e.,

$$\mathfrak{m}(x) = \bigoplus_{i=-1}^{-\mu} \mathfrak{g}_i(x), \quad \mathfrak{g}_i(x) := D^{-i}(x)/D^{-i-1}(x).$$

At each $x \in M$, the Lie bracket induces a bracket on $\mathfrak{m}(x)$ called the Levi bracket which is tensorial and turns $\mathfrak{m}(x)$ into a graded nilpotent Lie algebra (GNLA). The Lie algebra $\mathfrak{m}(x)$ is called the *symbol algebra* of $D(x)$. A distribution D of *constant type* has the property that $\mathfrak{m}(x) \cong \mathfrak{m}$ for some GNLA \mathfrak{m} for all $x \in M$.

An interesting class of GNLA arises from pairs (G, P) where G is a real or complex semi-simple group and $P \subset G$ is a parabolic subgroup in the following way. A $|\mu|$ -grading for the Lie algebra of G , denoted by \mathfrak{g} , is a vector space decomposition

$$\mathfrak{g} = \mathfrak{g}_\mu \oplus \dots \oplus \mathfrak{g}_\mu,$$

satisfying (i) $[\mathfrak{g}_i, \mathfrak{g}_j] \subset \mathfrak{g}_{i+j}$; (ii) $[\mathfrak{g}_i, \mathfrak{g}_{-1}] = \mathfrak{g}_{i-1}$ for all $i \leq -1$, i.e., the negative part of the grading,

$$\mathfrak{g}_- := \mathfrak{g}_{-\mu} \oplus \cdots \oplus \mathfrak{g}_{-1},$$

is generated by \mathfrak{g}_{-1} ; and (iii) $\mathfrak{g}_{\pm\mu} \neq 0$.

The subalgebra \mathfrak{g}_- is GNLA and

$$\mathfrak{p} := \mathfrak{g}_0 \oplus \mathfrak{g}_1 \oplus \cdots \oplus \mathfrak{g}_\mu$$

is always a parabolic subalgebra of \mathfrak{g} . Its corresponding subgroup, $P \subset G$, is a subgroup that lies between the normalizer $N_G(\mathfrak{p})$ and its identity component. The subgroup $G_0 \subset P$ defined as

$$G_0 = \{g \in P \mid \text{Ad}(\mathfrak{g}_i) \subset \mathfrak{g}_i \text{ for all } i = -\mu, \dots, \mu\}$$

is called the *Levi subgroup* corresponding to P .

For such a choice of (G, P) , there is a canonical construction of a bracket generating distribution whose symbol algebra is \mathfrak{g}_- . Let $N_{\mathfrak{g}}$ be the simply connected Lie group with Lie algebra \mathfrak{g}_- . The left-invariant vector fields obtained from \mathfrak{g}_{-1} induces a distribution of constant type on $N_{\mathfrak{g}}$ whose symbol algebra is isomorphic to \mathfrak{g}_- .

For a semi-simple Lie algebra \mathfrak{g} every derivation is inner. Therefore given a $|\mu|$ -grading of \mathfrak{g} it admits a unique *grading element* i.e., $e \in \mathfrak{g}_0$ such that $[e, g] = jg$ for all $g \in \mathfrak{g}_j$ and $-\mu \leq j \leq \mu$.

Consequently, the space of q -forms on \mathfrak{g}_- with values in \mathfrak{g} , denoted by $\Lambda^q(\mathfrak{g}_-, \mathfrak{g})$, admits a decomposition $\oplus_p \Lambda^{p,q}(\mathfrak{g}_-, \mathfrak{g})$ where $\Lambda^{p,q}(\mathfrak{g}_-, \mathfrak{g})$ is the set of q -forms ω , with weight p , i.e., $\mathcal{L}_e \omega = p\omega$.¹⁶ The decomposition can be written as

$$\Lambda^{p,q}(\mathfrak{g}_-, \mathfrak{g}) = \bigoplus_k \mathfrak{g}_{k+p} \otimes \wedge_k^q(\mathfrak{g}_-^*), \quad \text{where} \quad \wedge_k^q(\mathfrak{g}_-^*) := \sum_{\substack{r_1 + \dots + r_q = k \\ r_1, \dots, r_q < 0}} \mathfrak{g}_{r_1}^* \wedge \cdots \wedge \mathfrak{g}_{r_q}^*.$$

According to a theorem of Tanaka [Yam93], if $H^{1,p}(\mathfrak{g}_-, \mathfrak{g}) = 0$, for all $p \geq 0$, then via a process called the *Tanaka prolongation* the Lie algebra of infinitesimal symmetries of a distribution with symbol algebra \mathfrak{g}_- is isomorphic to \mathfrak{g} , i.e., $\inf(D) = \mathfrak{g}$.

The above theorem can be applied to *regular infinitesimal flag structures* of type (G, P) , which is defined as follows. Consider a manifold M endowed with a bracket generating distribution of constant type whose symbol algebra is \mathfrak{m} . The graded frame bundle is a right principal bundle $\text{Aut}_{gr}(\mathfrak{m}) \rightarrow \mathcal{F}_{gr}(M) \rightarrow M$, where $\text{Aut}_{gr}(\mathfrak{m})$ is the group of automorphisms of \mathfrak{m} that preserve the grading.¹⁷ If the symbol algebra of the distribution is isomorphic to \mathfrak{g}_- and $\text{Aut}_{gr}(\mathfrak{m})$ is isomorphic¹⁸ to $\text{Ad}(G_0) \subset \text{Aut}_{gr}(\mathfrak{g}_-)$, where G_0 is the Levi subgroup of (G, P) and Ad is the adjoint action, then one says that the distribution induces a regular infinitesimal flag structure of type (G, P) .

By a result of Tanaka [Tan79], Morimoto [Mor93] and Čap-Schichl [ČS00], there is an equivalence of categories between *regular*, *normal* parabolic geometries and regular infinitesimal flag structures via Tanaka prolongation. Regularity and normality of a parabolic geometry (G, P) is defined in terms of its curvature function of its Cartan connection (\mathcal{P}, ω) as follows.

Therefore, at each point $s \in \mathcal{P}$, the curvature $\Omega = d\omega + [\omega, \omega]$ of the Cartan connection can be used to define the curvature function $K : \mathcal{P} \rightarrow \wedge^2 \mathfrak{g}_-^* \otimes \mathfrak{g}$ as

$$K(s)(X, Y) = \Omega(\omega^{-1}(X)(s), \omega^{-1}(Y)(s)), \quad X, Y \in \mathfrak{g}_-,$$

where the space \mathfrak{g}_-^* can be identified with $\mathfrak{g}_+ := \oplus_{i>0} \mathfrak{g}_i$ via the Killing form. The Spencer cohomology groups $H^{p,q} \subset \mathfrak{g} \otimes \wedge^q(\mathfrak{g}_-^*)$, are defined using the Kostant operators $\partial : \Lambda^{p,q} \rightarrow$

¹⁶In [Yam93], the space $\Lambda^{p,q}$ is referred to as $C^{p-q+1,q}$.

¹⁷In case M is endowed with additional structure, e.g., conformal structures or semi-Riemannian structures, this principal bundle can be reduced further.

¹⁸If there exists additional structure on the distribution then $\text{Aut}_{gr}(\mathfrak{m})$ reduces to G_0 .

$\Lambda^{p-1,q+1}$, and $\partial^* : \Lambda^{p,q} \rightarrow \Lambda^{p+1,q-1}$, given by

$$\begin{aligned} \partial c(X_1 \wedge \cdots \wedge X_{q+1}) &= \sum_i (-1)^{i+1} [X_i, c(X_1 \wedge \cdots \wedge \hat{X}_i \wedge \cdots \wedge X_{q+1})] \\ &\quad + \sum_{i < j} (-1)^{i+j} c([X_i, X_j] \wedge X_1 \wedge \cdots \wedge \hat{X}_i \wedge \cdots \wedge \hat{X}_j \wedge \cdots \wedge X_{q+1}) \\ \partial^* c(X_1 \wedge \cdots \wedge X_{q-1}) &= \sum_i [e_i^*, c(e_i \wedge X_1 \wedge \cdots \wedge X_{q-1})] \\ &\quad + \frac{1}{2} \sum_{i,j} (-1)^{j+1} c([e_i^*, X_j]_- \wedge e_i \wedge X_1 \wedge \cdots \wedge \hat{X}_j \wedge \cdots \wedge X_{q-1}), \end{aligned}$$

where $c \in \Lambda^{p,q}(\mathfrak{g})$ and $[e_i^*, X_j]_-$ denotes the \mathfrak{g}_- -component of $[e_i^*, X_j]$, with respect to the decomposition $\mathfrak{g} = \mathfrak{g}_- \oplus \mathfrak{p}$. The operators above result in the orthogonal decomposition

$$(2.44) \quad \Lambda^{p,q} = H^{p,q} + \square \Lambda^{p,q},$$

where the operator $\square := \partial^* \partial + \partial \partial^*$ is referred to as the Kostant Laplacian and $\square H^{p,q} = 0$.

The $\Lambda^{p,2}$ -component of the curvature function K is denoted by $K^{(p)}$. Consequently, K can be decomposed as

$$(2.45) \quad K = \sum_{i=-\mu+2}^{3\mu} K^{(i)},$$

where $K^{(p)} : \mathfrak{g}_i \times \mathfrak{g}_j \rightarrow \mathfrak{g}_{i+j+p}$ is the component of K with homogeneity p . Additionally, one can decompose the curvature function as

$$K = \sum_{i=-\mu}^{\mu} K_p,$$

where K_p takes value in \mathfrak{g}_p . The \mathfrak{g}_- -component of K , $\sum_{i=-\mu}^{-1} K_i$, is denoted by K_- . The components of the curvature lying in $H^{p,2}$ are called *harmonic curvature* which constitute the set of essential invariants.

A Cartan connection is called *torsion-free* if $K_- = 0$, *normal* if $\partial^* K = 0$, and *regular* if it is normal and $K^{(i)} = 0$, for all $i \leq 0$.

2.4.2. Characterizing causal structures by a bracket generating distribution. The fact that $[\varepsilon^{ab}]$ is non-degenerate allows one to give a characterization of causal structures in terms of a bracket generating distribution with a certain small growth vector obtained as follows.

Define the distributions Δ and D such that $(\Delta)^\perp = \mathcal{I}_{\text{bas}}$ and $D^\perp = \{\omega^0, \dots, \omega^{n-1}\}$. The rank of D is n and Δ is a completely integrable sub-distribution of D of corank 1 whose integral manifolds are given by the fibers \mathcal{C}_x . Let \mathbf{v} be a characteristic vector field. It follows that $\mathbf{v} \in D$ and $\mathbf{v} \notin \Delta$.

Knowing $\omega^a \equiv -\varepsilon^{ab} \theta_b \wedge \omega^n$, modulo $I\{\omega^0, \dots, \omega^{n-1}\}$ and that Δ is completely integrable implies that $(D^2)^\perp = \{\omega^0\}$. Similarly, the equation $d\omega^0 \equiv -\theta_a \wedge \omega^a$, modulo $I\{\omega^0\}$, implies $(D^3)^\perp = \{0\}$, i.e. $D^3 = T\mathcal{C}$. As a result, the small growth vector of D is $(n, 2n-1, 2n)$.

Let $\{f_1, \dots, f_{n-1}\}$ be a dual basis for Δ and e_n be a basis for D/Δ . Since the growth vector is $(n, 2n-1, 2n)$, it follows that $\text{rank}(D^2/D) = n-1$. Because Δ is Frobenius, one obtains that the set of vectors $e_a = [e_n, f_a]$ are linearly independent and span D^2/D . This relation among e_a, e_n and f_a is equivalent to the structure equation $d\omega^a \equiv -\varepsilon^{ab} \theta_b \wedge \omega^n$ modulo $I\{\omega^0, \dots, \omega^{n-1}\}$.

Furthermore, the equation $d\omega^0 \equiv \theta_a \wedge \omega^a$ modulo $I\{\omega^0\}$, implies that there is a non-degenerate pairing between e_a 's and f_a 's, i.e., $[e_a, f_a] = e_0$ (no summation involved) for all $1 \leq a \leq n-1$, where e_0 spans D^3/D^2 .

One can investigate the action induced by the structure group G_5 in (2.34) on the *symbol algebra* of D as follows. Each tangent space $T_x \mathcal{C}$ admits a filtration $D(x) \subset D^2(x) \subset D^3(x)$ which results in the grading $\mathfrak{m}(x) = \mathfrak{g}_{-1}(x) \oplus \mathfrak{g}_{-2}(x) \oplus \mathfrak{g}_{-3}(x)$ where $\mathfrak{g}_i(x) = D^{-i}/D^{-i-1}$ with $D^0 = \{0\}$. Using the Lie bracket, $\mathfrak{m}(x)$ becomes a GNLA which would be the symbol algebra of D . Since \mathfrak{g}_{-1} is given by the kernel of $\mathcal{I}_1 := I\{\omega^0, \dots, \omega^{n-1}\}$, the action of the structure group G_5 on $\mathcal{I}_{\text{tot}} = I\{\omega^0, \dots, \omega^n, \dots, \theta^{n-1}\}$, modulo \mathcal{I}_1 , is given by scaling on ω^n and by conformal transformations on θ^a , i.e. ,

$$\omega^n \rightarrow \mathbf{b}^2 \omega^n, \quad \theta^a \rightarrow \frac{\mathbf{a}}{\mathbf{b}} \mathbf{A}^a_b \theta^b.$$

This action is isomorphic to the action of the *Levi subgroup* of $O(p+2, q+2)$ corresponding to the parabolic subgroup $P_{1,2}$ if $n \geq 3$ or $P_{1,2,3}$ if $n = 3$, on the coframe bundle. In other words, in causal structures the group of automorphisms of the coframe bundle which preserves the grading of $T_x\mathcal{C}$ is reduced to G_0 .

2.4.3. Normal Cartan connection. Using the flat model 2.2, it follows that the symbol algebra arising from the bracket generating distribution D , defined in Section 2.4.2 as $\{\omega^0, \dots, \omega^{n-1}\}^\perp$, is isomorphic to the negative part of the $|3|$ -grading for $\mathfrak{g} = \mathfrak{o}(p+2, q+2)$, i.e.,

$$\mathfrak{m} \cong \mathfrak{g}_- = \mathfrak{g}_{-3} \oplus \mathfrak{g}_{-2} \oplus \mathfrak{g}_{-1}.$$

The parabolic subalgebra that corresponds to this $|3|$ -grading is denoted as $\mathfrak{p}_{1,2}$ when $n \geq 4$ and $\mathfrak{p}_{1,2,3}$ when $n = 3$. The corresponding parabolic subgroup is given in Section 2.2, i.e., the parabolic subgroup of $O(p+2, q+2)$ that is isomorphic to the stabilizer of the subspace spanned by $\{e_1, e_2\}$ where $\{e_1, \dots, e_{n+3}\}$ is some basis for \mathbb{R}^{n+3} and $p+q = n-1$. Alternatively, $\mathfrak{p}_{1,2}$ can be described by crossing the first two nodes in the Dynkin diagrams B_n and D_n when $n \geq 4$ and all three nodes when $n = 3$.

Furthermore, the distribution D induces a regular infinitesimal flag structure. This is due to the fact that the condition $H^{p,1}(\mathfrak{g}_-, \mathfrak{g}) = 0$ is satisfied for all $p \geq 0$. More directly, by the discussion in Section 2.4.2 and the first order structure equations, one obtains that $\text{Aut}_{gr}(\mathfrak{g}_-)$ is reduced to the Levi subgroup. Thus, by Tanaka prolongation one can view causal structures as regular, normal parabolic geometries of type $(B_n, P_{1,2})$ or $(D_n, P_{1,2})$ if $n \geq 4$, or $(D_3, P_{1,2,3})$. In Section 2.5.1 it is discussed that when $n \geq 4$, causal structures are canonically isomorphic to regular, normal parabolic geometries of type $(B_n, P_{1,2})$ or $(D_n, P_{1,2})$. When $n = 3$, causal structures are “*subgeometries*” of regular, normal parabolic geometries of type $(D_3, P_{1,2,3})$, i.e., the integrability of the fibers \mathcal{C}_x , implies the vanishing of one of the harmonic curvature components of such parabolic geometries.

Knowing the existence of a normal Cartan connection for causal structures, the normality conditions of Tanaka are straightforward to obtain and can be used to recursively define the correct change of basis which transforms the $\{e\}$ -structure to a Cartan connection (see [Mak16]). Since finding the explicit expressions for the change of basis is very tedious and the result is not particularly illuminating, finding the normal Cartan connection is not carried out here.

2.5. Essential invariants and the Raychaudhuri equation. In this section the essential invariants whose vanishing imply the flatness of the causal structure are introduced. Their geometric interpretation is given. A variational problem is naturally associated to a causal geometry which allows one to define the notion of null Jacobi fields in the causal setting. Finally, the section ends with a discussion on associated Weyl connections of a causal structure and a derivation of the Raychaudhuri equation for causal structures. In this section there are references to the Appendix, particularly to the $\{e\}$ -structure obtained in Section A.1.4.

2.5.1. The harmonic curvature. According to the $\{e\}$ -structure obtained in Section A.1.4, obstructions to the flatness of a causal structure, as listed in Sections A.2 and A.3, are generated by F_{abc} and W_{anbn} . For instance, differentiating (2.37d) gives

$$(2.46) \quad \begin{aligned} W_{anbc} = & \frac{2}{3} (W_{an[b|n|;\underline{c}]} + F_{a[c}{}^d W_{b]ndn}) \\ & + \frac{4}{3(n+2)} (\varepsilon_{a[c} W_{n|b]n;\underline{d}} + \varepsilon_{a[b} F^{de}{}_{c]} W_{dnen}) \end{aligned}$$

In Section A.2, the quantities W_{ijkl} are expressed as the coframe derivatives of W_{anbn} with respect to $\frac{\partial}{\partial \theta^a}$. As a result, in the case $F_{abc} = 0$, the vanishing of W_{anbn} implies flatness of the causal structure.

For an understanding of W_{anbn} , consider the 2-plane given by $\lambda \wedge \mu$ where $\lambda = \lambda^i \mathbf{e}_i, \mu = \mu^i \mathbf{e}_i \in TM$ for some basis $\{\mathbf{e}_0, \dots, \mathbf{e}_n\}$. The *sectional conformal curvature* of a pseudo-conformal structure evaluated at $\lambda \wedge \mu$ is defined by

$$W(\lambda \wedge \mu) = W_{ijkl} \lambda^i \mu^j \lambda^k \mu^l.$$

As a result, at the point $(x, [y]) \in \mathcal{C}$, the invariant W_{anbn} can be interpreted as the part of the conformal sectional curvature acting on 2-planes $\xi^a = \frac{\partial}{\partial \omega^n} \wedge \frac{\partial}{\partial \omega^a}$. According to

the construction of the coframes in (2.11), ξ is a subset of $\widehat{T}_y\mathcal{C}_x$ and contains \hat{y} where $[\hat{y}] = y \in \mathcal{C}_x$. It follows that $\xi^a = \mu^a \wedge y^i \partial_i$, where μ^a is an element of the *shadow space*

$$(2.47) \quad S_{(x,[y])} := \widehat{T}_y\mathcal{C}_x / \hat{y}.$$

As a result, W_{anbn} can be thought of as the restriction of the bilinear form $W_{ijkl}y^jy^l$ to the shadow space. This restriction is well-defined up to scale. Infinitesimally, this scaling action is realized by the 1-form ϕ_n as is apparent from (2.39d).

Finally, it can be argued that the Fubini cubic form and the Wsf curvature coincide with the harmonic curvatures of the parabolic geometries that correspond to causal structures. By the discussion in Section 2.4.3, causal structures on an $(n+1)$ -dimensional manifold correspond to parabolic geometries of type $(B_n, P_{1,2})$ and $(D_n, P_{1,2})$ if $n \geq 4$, and $(D_3, P_{1,2,3})$. As discussed in Section 2.4.1, a grading is induced on the cohomology classes of parabolic geometries. Moreover, the set of essential invariants for a parabolic geometry is given by the harmonic curvature. It turns out that for parabolic geometries of type $(B_n, P_{1,2})$ and $(D_n, P_{1,2})$ if $n \geq 4$, the harmonic curvature has two irreducible components. Using the grading (2.45) of the curvature, one component has homogeneity one whose \mathfrak{g}_0 -irreducible representation is given as a totally symmetric and trace-free cubic tensor. The other irreducible component has homogeneity two whose \mathfrak{g}_0 -irreducible representation is given as a symmetric and trace-free bilinear form (see [KT14, ANN15, Yam93]). The structure equations of causal geometries imply that the essential invariant F_{abc} corresponds to a cohomology class of homogeneity one and the Wsf curvature corresponds to a cohomology class of homogeneity two. Also, from the infinitesimal group actions (2.39a) and (2.39d) one obtains that the Fubini cubic form and the Wsf curvature are closed under the action of the Levi subgroup. Moreover, one can impose the Tanaka normality conditions on the pseudo-connection form of the $\{e\}$ -structure to derive a normal Cartan connection (see [Mak16]). It turns out that the necessary change of coframe does not affect the Fubini cubic form and the Wsf curvature, i.e., the essential invariants of the $\{e\}$ -structure coincide with the harmonic curvature.

The situation is different in four dimensions. Four dimensional causal structures are a special class of parabolic geometries of type $(D_3, P_{1,2,3})$. The harmonic curvature of these parabolic geometries is five dimensional, three of which have homogeneity one and two have homogeneity two. Consider a four dimensional causal structure of split signature. The Fubini cubic form of projective surfaces and the Wsf curvature are comprised of two entries (see Section 4). By Equation (4.8) all these four entries are irreducible under the action of the Levi subgroup. As a result, four dimensional causal structures are special cases of parabolic geometries of type $(D_3, P_{1,2,3})$ in which one of the harmonic curvatures of homogeneity one is zero.

2.5.2. The ssf curvature. The action of the structure group G_5 defined in (2.34) on the bilinear form (2.35) is a scaling action $g \rightarrow \mathbf{a}^2 \mathbf{b}^2 g$. Fix a representative g and consider the sub-bundle $\mathcal{P}_1 = \{p \in \mathcal{P} \mid \mathbf{a}^2 \mathbf{b}^2 = 1\}$, where \mathcal{P} denotes the structure bundle associated to the derived $\{e\}$ -structure (see Section A.1.4). It follows that over \mathcal{P}_1 one has

$$(2.48) \quad 0 = d(\mathbf{a}^2 \mathbf{b}^2) = 2(\phi_0 + \phi_n) + s_i \omega^i + s_{\underline{a}} \theta^{\underline{a}},$$

for some functions s_i and $s_{\underline{a}}$. Differentiating this equation again, it follows that

$$(2.49) \quad ds_i \equiv \pi_i + \delta_i^0 (s_a \gamma^a + s_a \pi^a) + \delta_i^a (s_b \phi^b{}_a + s_n \gamma_a + s_{\underline{a}} \pi_n) \mod \mathcal{I}_{\text{tot}}.$$

Consequently, s_i can be translated to zero by the action of the structure group. It follows that

$$(2.50) \quad \pi_i = p_{ij} \omega^j + p_{i\underline{a}} \theta^{\underline{a}}.$$

Substituting (2.50) and $\phi_n = -\phi_0 + s_{\underline{a}} \theta^{\underline{a}}$ into the structure equations, equation (2.37d) becomes

$$(2.51) \quad d\theta^a = \phi^a{}_b \wedge \theta^b - 2\phi_0 \wedge \theta^a + \frac{1}{2} R^a{}_{nij} \omega^i \wedge \omega^j + E^a{}_{nbi} \theta^b \wedge \omega^i + \frac{1}{2} C^a{}_{nbc} \theta^b \wedge \theta^c.$$

where

$$(2.52) \quad \begin{aligned} R^a{}_{nbn} &= W^a{}_{nbn} + p_{nn} \delta^a{}_b, & R^a{}_{nbc} &= W^a{}_{nbc} + \delta^a{}_c p_{nb} + \delta^a{}_b p_{nc}, \\ R^a{}_{n0b} &= -p^a{}_b + \delta^a{}_b p_{n0}, & R^a{}_{n0n} &= p^a{}_n \end{aligned}$$

In conformal geometry, the symmetric part of p_{ij} is referred to as the *Schouten tensor*. The quantities R^a_{nbn} and p_{nn} are called the *shadow flag curvature* and the *scalar shadow flag curvature* (or the SSF curvature) of the causal structure respectively.

2.5.3. Null Jacobi fields and the tidal force. As was mentioned in section 2.3.5, the characteristic curves of causal structures is intrinsically defined by the quasi-contact structure of \mathcal{C} . Nevertheless, one can associate a *variational problem* to causal geometries via the Griffiths formalism [Gri83] (see also [Hsu92]). As a result, the notion of null Jacobi fields naturally arises. The corresponding Jacobi equations result in another interpretation of the WSF curvature. The fact that Jacobi equations of null Jacobi fields involve the WSF curvature has to do with the second Lie derivative of g in (2.35), which is referred to as the *tidal force* [O’N83].

Recall that for the Pfaffian system $\mathcal{I}_{\text{char}} = \{\omega^0, \dots, \omega^{n-1}, \theta^1, \dots, \theta^{n-1}\}$, after fixing an interval $I = (a, b)$, one can define the space of smooth immersions of its integral curves as

$$\widehat{\mathcal{W}}(I) = \{\gamma : I \rightarrow \mathcal{C} \mid \gamma^*(\mathcal{I}_{\text{tot}}) = 0\}.$$

The space $\widehat{\mathcal{W}}(I)$ can be given a Whitney C^∞ -topology. After making a choice of orientation for the characteristic curves, the space of the characteristic curves of \mathcal{C} , $\mathcal{W}(I)$, is defined as $\widehat{\mathcal{W}}(I)/\text{Diff}(I)$, i.e., the space obtained from $\widehat{\mathcal{W}}(I)$ by identifying the characteristic curves that are reparametrizations of each other.

Consider the functional $\Phi : \mathcal{W} \rightarrow \mathbb{R}$, which at a point $[\gamma] \in \mathcal{W}(I)$ is defined by

$$\Phi([\gamma]) = \int_I \gamma^* \omega^0.$$

The triple $(\mathcal{C}, I, \omega^0)$ is referred to as a *variational problem* on \mathcal{C} , and is intrinsic to any causal structure. Since $\omega^0(\dot{\gamma}) = 0$ everywhere, the above definition of Φ is independent of the choice of representative $\gamma \in [\gamma]$. This is in contrast with Finsler geometry where the geodesics are defined as the extremal curves of the arc-length functional which is always positive and are equipped with a specific parametrization.

Intuitively, the infinitesimal variations of a characteristic curve γ can be identified with the tangent space $T_\gamma \mathcal{W}(I)$ which is the space of smooth variational vector fields along γ . To give a more precise definition consider a parametrization for $I = (a, b)$ and a map $\Gamma : (-\epsilon, \epsilon) \rightarrow \mathcal{W}(I)$. As a result of such parametrization, the curves given by $\Gamma(s)$ can be parametrized. Using this parametrization, one can define $\widehat{\Gamma} : (-\epsilon, \epsilon) \times I \rightarrow \mathcal{C}$, as $\widehat{\Gamma}(s, t) = (\Gamma(s))(t)$. The map Γ is called a compact variation of the curve γ if the corresponding map $\widehat{\Gamma}$ is smooth, $\widehat{\Gamma}(0, t) = \gamma(t)$, $\widehat{\Gamma}^*(s_0, t)(\mathcal{I}_{\text{char}}) = 0$, for any fixed s_0 , and $\widehat{\Gamma}$ is compactly supported in I , i.e., for all values of s , the curves $\widehat{\Gamma}(s, t)$ and $\widehat{\Gamma}(0, t)$ coincide outside of a compact subset of I . Then, $T_\gamma \mathcal{W}(I)$ is expressed as

$$T_\gamma \mathcal{W}(I) := \left\{ \frac{\partial \widehat{\Gamma}}{\partial s}(0, t) \mid \Gamma : (-\epsilon, \epsilon) \rightarrow \mathcal{W}(I) \text{ is a compact variation of } \gamma \right\}.$$

The *variational equations* of a characteristic curve can be thought of as the first order approximation of $T_\gamma \mathcal{W}(I)$. A practical derivation of the variational equations can be given in terms of the Pfaffian system $\mathcal{I}_{\text{char}}$.

Let $\eta \in \mathcal{I}_{\text{char}}$, and Γ a variation of the characteristic curve γ . By the definition of $\widehat{\Gamma}(s, t)$, one has $\widehat{\Gamma}^*(s_0, t)\eta = 0$, and therefore $\mathcal{L}_{\frac{\partial}{\partial s}} \widehat{\Gamma}^*(s, t)\eta \equiv 0$ modulo $I\{ds\}$. Setting $s = 0$, the latter expression implies that

$$\gamma^*(\hat{J} \lrcorner d\eta + d(\hat{J} \lrcorner \eta)) = 0,$$

where $\hat{J}(t) = \frac{\partial}{\partial s} \widehat{\Gamma}(0, t)$ is the infinitesimal variations associated to $\Gamma(s)$.

In the expression of the variational equations one can drop the pull-back γ^* and write them as

$$\hat{J} \lrcorner d\eta + d(\hat{J} \lrcorner \eta) \equiv 0 \quad \text{mod} \quad \mathcal{I}_{\text{char}}.$$

The tangential component of the vector field $\hat{J}(t)$ along $\gamma(t)$ has no effect on the variational equations. Therefore, \hat{J} along the curve $\gamma(t)$ can be expressed as

$$\hat{J}(t) = V^a(t) \frac{\partial}{\partial \omega^a} + V^a(t) \frac{\partial}{\partial \theta^a}.$$

The vector field $J = \pi_* \hat{J}$ is called a *null Jacobi field* along the null geodesics $\pi(\gamma(t))$.

The variational equations for $\hat{J}(t)$ can be written as

$$(2.53a) \quad \hat{J} \lrcorner d\omega^0 + d(\hat{J} \lrcorner \omega^0) = 0,$$

$$(2.53b) \quad \hat{J} \lrcorner d\omega^a + d(\hat{J} \lrcorner \omega^a) = 0, \quad \text{mod } I\{\omega^0, \dots, \omega^n, \theta_1, \dots, \theta_{n-1}\}.$$

$$(2.53c) \quad \hat{J} \lrcorner d\theta_a + d(\hat{J} \lrcorner \theta_a) = 0,$$

Equation (2.53a) gives no new information. The last two equations give

$$(2.54) \quad \begin{aligned} dV^a + V^b \phi^a_b - V^a \omega^n &\equiv 0, \\ dV^a + V^b \phi^a_b + W^a_{bn} V^b \omega^n &\equiv 0, \quad \text{mod } \mathcal{I}_{\text{char}}. \end{aligned}$$

Using the pseudo-connection from Section 2.3.12, the covariant derivative along characteristic curves of vector fields lying within the shadow space (2.47) is given by

$$D_{\mathbf{v}} \hat{J} \equiv \mathbf{v} \lrcorner \left(dV^a \frac{\partial}{\partial \omega^a} + V^b \phi^a_b \frac{\partial}{\partial \omega^a} \right), \quad \text{mod } \frac{\partial}{\partial \theta^1}, \dots, \frac{\partial}{\partial \theta^{n-1}},$$

where \mathbf{v} is the characteristic vector field satisfying $\omega^n(\mathbf{v}) = 1$.

Using equations (2.54), it follows that

$$(2.55) \quad \begin{aligned} D_{\mathbf{v}} D_{\mathbf{v}} \hat{J} &\equiv -W^a_{bn} V^b \frac{\partial}{\partial \omega^a}, \quad \text{mod } \frac{\partial}{\partial \theta^1}, \dots, \frac{\partial}{\partial \theta^{n-1}} \\ &\equiv -\overset{\text{sf}}{W} \hat{J}, \quad \text{mod } \frac{\partial}{\partial \theta^1}, \dots, \frac{\partial}{\partial \theta^{n-1}} \end{aligned}$$

where $\overset{\text{sf}}{W}$ is viewed as an automorphism of the shadow space. Recall from Section 2.5.1 that W^a_{bn} is given by $\omega^a(W(\frac{\partial}{\partial \omega^b}, \mathbf{v})\mathbf{v})$. Hence, defining

$$J' = \pi_*(D_{\mathbf{v}} \hat{J}),$$

the null Jacobi equation can be expressed as

$$(2.56) \quad J'' + W(J, \mathbf{v})\mathbf{v} = 0,$$

where $J = \pi_* \hat{J}$ and $W(J, \mathbf{v})\mathbf{v}$ denotes $\pi_*(W(\hat{J}, \mathbf{v})\mathbf{v})$. For convenience, in Section 2.5.4 Equation (2.56) is expressed as

$$(2.57) \quad J'' + \overset{\text{sf}}{W} J = 0.$$

In some references [O'N83] (cf. [HS11]) the automorphism of the shadow space $W^a_{nba} : S_{(x, [y])} \rightarrow S_{(x, [y])}$ is called the trace-free part of the *tidal force* in analogy with the terminology from General Relativity. Equation (2.55) shows that the WSF curvature plays a rule analogous to that of the conformal sectional curvature in the case of variational theory of null geodesics.

The relation between the tidal force and the trace-free part of the WSF curvature can be realized differently via the second Lie derivative the bilinear form g in (2.35) along the characteristic curves (cf. [HS11, Bry02]). The structure equations (2.37) imply

$$\mathcal{L}_{\mathbf{v}}^2 g = 4\omega^0 \circ \pi_n - 2\varepsilon_{ab} \theta^a \circ \theta^b - 2W_{anbn} \omega^a \circ \omega^b.$$

Therefore, for any two vector fields $X_1, X_2 \in \text{Ker } \omega^0$, $X_\epsilon = X_\epsilon^a \frac{\partial}{\partial \omega^a} + X_\epsilon^a \frac{\partial}{\partial \theta^a}$ where $\epsilon = 1, 2$, it follows that

$$W_{anbn} X_1^a X_2^b = -\frac{1}{2} \mathcal{L}_{\mathbf{v}}^2 g(X_1, X_2) - \varepsilon_{ab} X_1^a X_2^b.$$

2.5.4. The Raychaudhuri equation. In this section, a causal analogue of the Raychaudhuri equation along null geodesics is derived. The approach that is taken here is presented in [BEE96, Chapter 12].

Taking a section of the structure bundle, $s : \mathcal{C} \rightarrow \mathcal{P}$, define the null Jacobi fields $\hat{J}_1(t), \dots, \hat{J}_{n-1}(t)$ along the characteristic curve $\gamma(t)$, $t \in [a, b]$ by the initial conditions

$$\hat{J}_a(t_0) = 0, \quad D_{\mathbf{v}(t_0)} \hat{J}_a(t_0) = \frac{\partial}{\partial (s^* \omega^a)}$$

where $t_0 \in [a, b]$ and $\mathbf{v}(t)$ is tangent vector to $\gamma(t)$. Define the $(1, 1)$ tensor field called the *null Jacobi tensor* as

$$A(t) = [J_1(t), \dots, J_{n-1}(t)],$$

which is a $(n-1) \times (n-1)$ matrix whose b th column is the null Jacobi field $J_b(t) = \pi_*(\hat{J}_b(t))$.

The optical invariants along null geodesics are defined as follows.

Definition 2.17. Let A be a null Jacobi tensor along a null geodesic and define $B = A'A^{-1}$, where A is invertible. The *expansion*, θ , *vorticity tensor*, ω and *shear tensor*, σ , are defined as

$$\theta = B^a_a, \quad \omega^a_b = \frac{1}{2} (B^a_b - (B^*)^a_b), \quad \sigma = \frac{1}{2} (B^a_b + (B^*)^a_b) - \frac{\theta}{n-1} \delta^a_b,$$

where B^* denotes the adjoint matrix with respect to the bilinear form g .

Following the discussion in Section 2.5.3, after taking a section of the structure bundle one obtains that a null Jacobi field satisfies the equation

$$J'' + R^{\text{sf}} J = 0,$$

rather than Equation (2.57), where R^{sf} represents the matrix form of the shadow flag curvature $R^a_{\text{nb}n} = W^a_{\text{nb}n} + p_{nn} \delta^a_b$ defined in (2.52).

It follows that

$$A'' + R^{\text{sf}} A = 0.$$

Moreover, one has

$$B' = (A'A^{-1})' = A''A^{-1} - A'A^{-1}A'A^{-1} = -R^{\text{sf}} - B^2.$$

Since $B = \omega + \sigma + \frac{\theta}{n-1} \text{Id}$, one obtains

$$\theta' = \text{tr}(B') = -\text{tr}(R^{\text{sf}}) - \text{tr}(B^2) = -\text{tr}(R^{\text{sf}}) - \text{tr} \left[\left(\omega + \sigma + \frac{\theta}{n-1} \text{Id} \right)^2 \right]$$

Using the fact that $\text{tr}(\omega) = \text{tr}(\sigma) = \text{tr}(\omega\sigma) = 0$ and $\text{tr}(R^{\text{sf}}) = (n-1)p_{nn}$ one arrives at

$$(2.58) \quad \theta' = -(n-1)p_{nn} - \text{tr}(\omega^2) - \text{tr}(\sigma^2) - \frac{\theta^2}{n-1},$$

which is the causal analogue of the Raychaudhuri equation along null geodesics.

2.6. Examples. In this section three examples of causal structures are given. The first one is obtained from the product of two Finsler structures and the second and third ones are given in terms of a defining function. These examples are used in Section 4.

2.6.1. Product of Finsler structures. A Finsler manifold is denoted by (N, Σ) , where N is n -dimensional and $\Sigma \subset TN$ is of codimension one whose fibers $\Sigma_p \subset T_p N$ are strictly convex hypersurfaces. The fiber bundle Σ is often referred to as the indicatrix bundle over which the Finsler metric has unit value. It is known [Car34, Bry02] that Σ is equipped with unique coframing $\eta = (\eta^0, \eta^1, \dots, \eta^{n-1}, \zeta^1, \dots, \zeta^{n-1})$ satisfying

$$(2.59) \quad \begin{aligned} d\eta^0 &= -\zeta_a \wedge \eta^a, \\ d\eta^a &= \zeta^a \wedge \eta^0 - \psi^a_b \wedge \eta^b - I^{ab}_c \zeta_b \wedge \eta^c, \\ d\zeta_a &= \psi^b_a \wedge \zeta_b + R_{0a0b} \eta^0 \wedge \eta^b + \frac{1}{2} R_{0abc} \eta^b \wedge \eta^c - J^b_a \zeta_b \wedge \eta^c \end{aligned}$$

for some invariants I_{abc} , J_{abc} , R_{0a0b} and R_{0abc} . The 1-form η^0 is called the Hilbert form of the Finsler structure which, at the point $v \in \Sigma_p$, is uniquely characterized by the conditions

$$\eta^0(H_v \Sigma_p) = 0, \quad \eta^0(v) = 1,$$

where $H_v \Sigma_p \subset T_p N$ is the tangent hyperplane to Σ_p at the point v . As a result of the structure equation, η^0 induces a contact structure on Σ whose associated Reeb vector field defines the geodesic flow of the Finsler structure on Σ . Moreover, the quadratic form $\omega^0 \circ \omega^0 + \delta_{ab} \omega^b \circ \omega^c$ is well-defined for the Finsler structure. It turns out that the quantity $I_{abc} = \delta_{ad} \delta_{be} I^{de}_c$ can be used to define the cubic form $I_{abc} \zeta^a \circ \zeta^b \circ \zeta^c$, which, when restricted to a fiber Σ_p , coincides with the centro-affine cubic form of the hypersurface $\Sigma_p \subset T_p N$. The quantities J_{abc} measure the rate of change of I_{abc} along the geodesics, i.e., $J_{abc} = \frac{\partial}{\partial \eta^0} I_{abc}$. The quantity R_{0a0b} is symmetric using which the *flag curvature* of the Finsler structure can be represented by

$$R_{0a0b} \eta^a \circ \eta^b.$$

Let (N, Σ) and $(\tilde{N}, \tilde{\Sigma})$ be two Finsler manifolds. Given a strictly positive function f on their product space, at a point $p \in M = N \times_f \tilde{N}$, one can define a projective hypersurface whose associated cone is given by

$$(2.60) \quad \widehat{\mathcal{C}}_p = \{(v, \tilde{v}) \in T_p(N \times \tilde{N}) \mid \lambda f v \in \Sigma_{\pi(p)}, \lambda \tilde{v} \in \tilde{\Sigma}_{\tilde{\pi}(p)} \text{ for some } \lambda \in \mathbb{R}\}.$$

As a result, M is equipped with a causal structure of signature (n, \tilde{n}) where $\tilde{n} = \dim \tilde{N}$. Note that it is assumed that the Finsler structures are reversible.

If G and \tilde{G} represent the Finsler metrics on N and \tilde{N} , respectively, then the causal structure on M is locally given as the zero locus of $G - f\tilde{G}$. If f only depends on N , then the above causal structure corresponds to the warped product of the Finsler structures on N and \tilde{N} .

More specifically, let $M = N \times_f \mathbb{R}$, where N is a 3-dimensional Finsler manifold. On N there is the unique coframing $(\eta^0, \eta^1, \eta^2, \zeta^1, \zeta^2)$ satisfying (2.59). If η^3 represents a 1-form on \mathbb{R} , one has

$$df = f_{,0}\eta^0 + f_{,1}\eta^1 + f_{,2}\eta^2 + f_{,3}\eta^3,$$

where $f_{,i} = \frac{\partial}{\partial \eta^i} f$ are called the coframe derivatives of f . The projective Hilbert form is a multiple of

$$\omega^0 = \frac{1}{\sqrt{2}}(\eta^0 - f\eta^3),$$

where, by abuse of notation, the pull-backs are dropped and η^0 and η^3 represent $\pi^*(\eta^0)$ and $\tilde{\pi}^*(\eta^3)$.

A choice of 3-adapted coframing for the causal structure on M is given by

$$(2.61) \quad \begin{aligned} \omega^1 &= \eta^1, & \omega^2 &= \eta^2, & \omega^3 &= \frac{1}{\sqrt{2}}(-\eta^0 - f\eta^3), \\ \theta_1 &= \frac{1}{\sqrt{2}}\zeta_1 - \frac{1}{\sqrt{2}}f_{,1}\eta^3, & \theta_2 &= \frac{1}{\sqrt{2}}\zeta_2 - \frac{1}{\sqrt{2}}f_{,2}\eta^3, \\ \gamma_1 &= -\frac{1}{\sqrt{2}}\zeta_1 + \frac{1}{\sqrt{2}}f_{,1}\eta^3, & \gamma_2 &= -\frac{1}{\sqrt{2}}\zeta_2 + \frac{1}{\sqrt{2}}f_{,2}\eta^3, \\ \phi_0 &= -\phi_3 = -\frac{1}{2}f_{,0}\eta^3, & \phi &= \pi_3 = 0. \end{aligned}$$

The quantities $F^{ab}{}_c$ in the structure equations (2.26) coincide with $I^{ab}{}_c$ in (2.59). By absorbing the trace of F_{abc} and K_{ab} , a 5-adapted coframing $(\tilde{\omega}^i, \tilde{\theta}_a)$ is given by

$$\tilde{\omega}^0 = \omega^0, \quad \tilde{\omega}^a = \omega^a + A^a \omega^0, \quad \tilde{\omega}^3 = \omega^3 + B\omega^0, \quad \tilde{\theta}_a = \theta_a,$$

where

$$\begin{aligned} A^1 &= -\frac{1}{2}(I_{111} + I_{122}), & A^2 &= \frac{1}{2}(I_{112} + I_{222}), \\ B &= \frac{1}{4}((A^1)^2 + (A^2)^2) - \frac{1}{2\sqrt{2}}(A^1{}_{,\underline{1}} + A^2{}_{,\underline{2}}) - A^1 F_{111} - A^2 F_{222}, \end{aligned}$$

where $F_{111} = \frac{1}{4}(I_{111} - 3I_{122})$, $F_{222} = \frac{1}{4}(I_{222} - 3I_{211})$ are the components of the Fubini cubic form of the causal structure and $A^a{}_{,\underline{b}}$ denotes the coframe derivative $\frac{\partial}{\partial \zeta^b} A^a$. A choice of connection forms for the first order structure equations is given by

$$\begin{aligned} \tilde{\gamma}^1 &= \gamma^1 + B_{,1}\omega^0 - \left(\frac{1}{\sqrt{2}}B_{,0} + A^1{}_{,1}\right)\tilde{\omega}^1 - \frac{1}{2}(A^1{}_{,2} + A^2{}_{,1})\tilde{\omega}^2 + \frac{1}{\sqrt{2}}A^1{}_{,0}\tilde{\omega}^3 + B\theta_1, \\ \tilde{\gamma}^2 &= \gamma^2 + B_{,2}\omega^0 - \frac{1}{2}(A^1{}_{,2} + A^2{}_{,1})\tilde{\omega}^1 - \left(\frac{1}{\sqrt{2}}B_{,0} + A^2{}_{,2}\right)\tilde{\omega}^2 + \frac{1}{\sqrt{2}}A^2{}_{,0}\tilde{\omega}^3 + B\theta_2, \\ \tilde{\phi}_0 &= -\frac{1}{2}A^1\theta_1 - \frac{1}{2}A^2\theta_2, \\ \tilde{\phi}_3 &= \frac{1}{2\sqrt{2}}(-B_{,0}\omega^0 + A^1{}_{,0}\tilde{\omega}^1 + A^2{}_{,0}\tilde{\omega}^2), \\ \tilde{\phi} &= \frac{1}{2}(A^1{}_{,2} - A^2{}_{,1})\omega^0 + \frac{1}{2\sqrt{2}}A^2{}_{,0}\tilde{\omega}^1 - \frac{1}{2\sqrt{2}}A^1{}_{,0}\tilde{\omega}^2 - \frac{1}{2}A^2\theta_1 + \frac{1}{2}A^1\theta_2 \\ \tilde{\pi}_3 &= -\frac{1}{\sqrt{2}}A^1{}_{,0}\theta_1 - \frac{1}{\sqrt{2}}A^2{}_{,0}\theta_2 - \frac{1}{2}(R_{0101} + R_{0202} + (\ln f)_{,11} + (\ln f)_{,22})\tilde{\omega}^3. \end{aligned}$$

Consequently, the WSF curvature is given by the trace-free part of

$$(2.62) \quad \frac{1}{2}R_{0a0b} + \frac{1}{2}(\ln f)_{,ab},$$

where $(\ln f)_{,ab} := \frac{\partial}{\partial \eta^b} \frac{\partial}{\partial \eta^a} \ln f$. Moreover, the quantities K_{ab} and L_a are given by

$$K_{12} = K_{21} = -\sqrt{2}A^1{}_{,\underline{2}} - F_{112}A^1 - F_{122}A^2, \quad K_{11} = -K_{22} = -\frac{1}{\sqrt{2}}(A^1{}_{,\underline{1}} - A^2{}_{,\underline{2}}), \quad L_a = -\sqrt{2}B_{,\underline{a}}.$$

To see the symmetric relation $K_{12} = K_{21}$ one needs to realize $A^1{}_{,\underline{2}} = A^2{}_{,\underline{1}}$. It is rather straightforward to generalize this example to obtain a causal structure on $M = N \times_f \mathbb{R}$

where N is an n -dimensional Finsler manifold. The corresponding WSF is given by expression (2.62) and the Fubini cubic form is obtained from the centro-affine cubic form of indicatrices of the Finsler structure.

2.6.2. *Cayley's cubic scroll.* Using the description (2.2), consider the causal structure on a 4-dimensional manifold given locally as the graph of the function

$$F(x^i, y^a) = \frac{1}{3}c_{111}(y^1)^3 + c_{12}y^1y^2 + c_1y^1 + c_2y^2 + c_0,$$

where the coefficients $c_{i\dots k}$ are functions of x^i 's.

A choice of 1-adapted coframing is given by

$$(2.63) \quad \begin{aligned} \omega^0 &= dx^0 - \frac{\partial}{\partial y^a} F dx^a + (y^a \frac{\partial}{\partial y^a} F - F) dx^3, \\ \omega^a &= dx^a - y^a dx^3, \\ \omega^3 &= dx^3, \\ \theta_a &= \frac{\partial^2}{\partial x^i \partial y^a} F dx^i - (\frac{\partial}{\partial x^a} F + \frac{\partial}{\partial x^0} F \frac{\partial}{\partial y^a} F) + \frac{\partial^2}{\partial y^a \partial y^b} F dy^b. \end{aligned}$$

A 2-adapted coframing is given by

$$\omega^1 \longrightarrow \omega^1 - \frac{c_{111}}{c_{12}} y^1 \omega^2, \quad \omega^3 \longrightarrow \frac{1}{c_{12}} \omega^3, \quad \theta_1 \longrightarrow \theta_1 - \frac{c_{111}}{c_{12}} y^1 \theta_2,$$

with respect to which the second fundamental form is normalized to $h^{ab} = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$. The details of higher coframe adaptations won't be discussed. This example is discussed further in Section 4.1.

It should be noted that when the only non-zero coefficients are $c_{12} = 1$ and $c_{00} = (x^1)^2$, then one obtains a local description of the split signature pp-wave metric. Moreover, when $c_0 = c_1 = c_2 = 0$ one obtains a local description of causal structures whose null cones are at each point projectively equivalent to Cayley's cubic scroll whose graph can be expressed as $y^0 = \frac{1}{3}(y^1)^3 + y^1 y^2$ in \mathbb{P}^3 . In Section 4.2, this example is used to characterize the submaximal causal structure of split signature.

2.6.3. *Locally isotrivially flat causal structures.* This class of causal structures are the analogue of locally Minkowskian Finsler structures.

Definition 2.18. A causal structure (\mathcal{C}, M^{n+1}) is called locally isotrivially flat if it is locally equivalent to $(M \times \mathcal{V}, M)$ where $\mathcal{V} \subset \mathbb{P}^n$ is a projective hypersurface.

Alternatively, a locally isotrivially flat causal structure has the property that locally it can be expressed as the graph of a function $F(x; y)$, as in (2.2), which has no dependency on x^i 's.

The local expressions (2.10) and (2.16) of the 1-forms ω^i and θ_a can be used to express the Fubini cubic form and WSF curvature. Firstly, note that $\theta_a = \partial_a \partial_b F dy^b$. To express F_{abc} , consider the following transformation

$$(2.64) \quad \tilde{\omega}^0 = \mathbf{a}^2 \omega^0, \quad \tilde{\omega}^a = \mathbf{E}^a \omega^0 + \omega^a, \quad \tilde{\omega}^n = \mathbf{e} \omega^0 + \omega^n, \quad \tilde{\theta}_a = \mathbf{a}^2 \theta_a.$$

Differentiating $\tilde{\omega}^0$, it follows that

$$d\tilde{\omega}^0 = -2\phi_0 \wedge \tilde{\omega}^0 - \tilde{\theta}_a \wedge \tilde{\omega}^a.$$

where $\phi_0 = -\frac{d\mathbf{a}}{\mathbf{a}} - \frac{\mathbf{E}^a}{2\mathbf{a}^2} \tilde{\theta}_a$. Consequently, in Equation (2.20b) for 1-adapted coframes one obtains

$$\gamma^a{}_0 = \frac{1}{\mathbf{a}^2} d\mathbf{E}^a + \frac{1}{\mathbf{a}^4} (\mathbf{E}^a \mathbf{E}^b + H^{ab}) \theta_b, \quad \psi^a{}_b = \frac{1}{\mathbf{a}^2} \mathbf{E}^a \theta_b, \quad h^{ab} = \frac{1}{\mathbf{a}^2} H^{ab},$$

where $[H^{ab}]$ denotes the inverse matrix of the vertical Hessian $[\partial_a \partial_b F]$.

By composing Equations (2.21) and (2.25), it follows that the Fubini cubic form F^{abc} is defined via the equation

$$dh^{ab} + h^{ac} \psi^b{}_c + h^{bc} \psi^a{}_c - 2h^{ab} (\phi_0 + \phi_n) \equiv F^{abc} \theta_c \mod \mathcal{I}_{\text{bas}}.$$

Transformations (2.64) imply that $\phi_n = 0$. Using the above expressions for ϕ_0 and $\psi^a{}_b$, without normalizing h^{ab} to ε^{ab} , one can impose the apolarity condition $h^{ab} F_{abc} = 0$, as in (2.30), by setting

$$\mathbf{a} = (\det(\partial_a \partial_b F))^{\frac{1}{2(n+1)}}, \quad \mathbf{E}^d = -\frac{\mathbf{a}^2}{n+1} H^{ab} H^{cd} \partial_a \partial_b \partial_c F.$$

Finally, using $\partial_a \partial_b F$ to lower indices, it follows that

$$(2.65) \quad F_{abc} = \mathbf{a}^{-2} \left(\partial_a \partial_b \partial_c F - \frac{1}{n+1} (\partial_a \partial_b F F_c + \partial_b \partial_c F F_a + \partial_c \partial_a F F_b) \right),$$

where $F_a = H^{bc} \partial_a \partial_b \partial_c F$.

The above computations go through for any causal structure. The expression (2.65) (see also [Sas99, AG93]) shows that the Fubini cubic form is a third order invariant. Using the Bianchi identities (A.19), it follows that the jet order of the invariants K^{ab} and L^a over \mathcal{C} are four and five, respectively.

To find the WSF curvature, note that without normalizing h^{ab} , the quantities E_{anb} in Equation (2.25) are defined via

$$dh^{ab} + h^{ac} \psi_c^b + h^{bc} \psi_c^a - 2h^{ab} (\phi_0 + \phi_n) = E_n^a{}^b \omega^n, \quad \text{mod } I\{\omega^0, \dots, \omega^{n-1}, \theta_1, \dots, \theta_{n-1}\},$$

which by isotrivally flatness is zero. Hence, the reduction of π_{ab} via (2.27), gives $\pi_{ab} - \frac{1}{n-1} \varepsilon^{cd} \pi_{cd} \varepsilon_{ab} = 0$. Combining this with the fact that $\theta_a = 0$ modulo $I\{\theta_1, \dots, \theta_{n-1}\}$, it follows that $W_{nbn}^a = 0$, i.e., locally isotrivally flat causal structures have vanishing WSF curvature.

3. CAUSAL STRUCTURES WITH VANISHING WSF CURVATURE

In this section causal structures with vanishing WSF curvature are studied. For such classes, a compatibility condition is derived which can be used to give a causal analogue of Landsberg spaces in Finsler geometry (see [BCS00] for an account of various types of Finsler structures). Moreover, under certain conditions, a rigidity theorem similar to that of Akbar-Zadeh's [AZ88] is obtained. Finally, the section ends with a discussion on Lie contact structures on the locally defined space of characteristic curves of a causal structure with vanishing WSF curvature.

3.1. A compatibility condition. In this Section a class of causal structures which can be regarded as analogues of Landsberg spaces in Finsler geometry is introduced. Their similarity is the content of Proposition 3.1.

Since DF_{abc} in (2.39a) (2.39a) is semi-basic,

$$(3.1) \quad dF_{abc} - F_{abd} \phi_c^d - F_{adc} \phi_b^d - F_{dbc} \phi_a^d - F_{abc} (\phi_0 - \phi_n) = F_{abc;i} \omega^i + F_{abc;\underline{a}} \theta^{\underline{a}}.$$

After taking the Lie derivative of (3.1) along characteristic curves and using the structure equations, one arrives at

$$df_{abc} - f_{abd} \phi_c^d - f_{adc} \phi_b^d - f_{dbc} \phi_a^d - f_{abc} (\phi_0 + \phi_n) + F_{abc} \pi_n \equiv 0 \quad \text{mod } \mathcal{I}_{\text{tot}}.$$

Assume $f_{abc} = \lambda F_{abc}$, for some function λ . Then, if $F_{abc} \neq 0$, the infinitesimal group action on λ is given by

$$d\lambda - 2\lambda \phi_n + \pi_n \equiv 0 \quad \text{mod } \mathcal{I}_{\text{tot}}.$$

By translating λ to zero, the 1-form π_n reduces to

$$(3.2) \quad \pi_n = p_{ni} \omega^i + p_{n\underline{a}} \theta^{\underline{a}}.$$

for some coefficients p_{ni} and $p_{n\underline{a}}$. Replacing π_n with the above expression in equation (2.37d), one finds the infinitesimal action of the group G_5 on $p_{n\underline{a}}$ to be given by

$$dp_{n\underline{a}} - p_{n\underline{a}} (\phi_0 + \phi_n) - p_{n\underline{d}} \phi_a^d + \pi_a \equiv 0, \quad \text{mod } \mathcal{I}_{\text{tot}}.$$

Translating $p_{n\underline{a}}$ to zero results in the reduction

$$(3.3) \quad \pi_a = p_{ai} \omega^i + p_{a\underline{b}} \theta^{\underline{b}}.$$

Consequently, equation (2.37d) changes to

$$(3.4) \quad d\theta_a = (\phi_a^b - (\phi_0 - \phi_n) \delta_a^b) \wedge \theta_b + \frac{1}{2} W_{anij} \omega^i \wedge \omega^j + E_{anb0} \theta^b \wedge \omega^0,$$

where

$$\begin{aligned} W_{anbn} &= p_{nn}, & W_{anbc} &= -p_{nb} \varepsilon_{ac} + p_{nc} \varepsilon_{ab}, \\ W_{an0n} &= p_{a0}, & W_{an0b} &= p_{ab}, & E_{anb0} &= -p_{a\underline{b}}. \end{aligned}$$

Similarly, one finds that the normalization $\varepsilon^{ab} p_{a\underline{b}} = 0$ results in $\pi_0 \equiv 0$ modulo \mathcal{I}_{tot} . It follows that

$$(3.5) \quad \pi_0 = p_{0i} \omega^i + p_{0\underline{a}} \theta^{\underline{a}}.$$

Assume furthermore that $p_{nn} \neq 0$. Using the relation (2.39d), similar to the derivation of (3.10), the infinitesimal group action on p_{nn} is found to be

$$(3.6) \quad dp_{nn} - 4p_{nn}\phi_n \equiv 0, \quad \text{mod } \mathcal{I}_{\text{tot}}.$$

Hence, the normalization $p_{nn} = \pm 1$ can be made, resulting in ϕ_n reduced to

$$(3.7) \quad \phi_n = r_i \omega^i + r_{\underline{a}} \theta^{\underline{a}}.$$

Furthermore, the infinitesimal group action of the reduced structure group on $r_{\underline{a}}$ is given by

$$dr_{\underline{a}} + r_{\underline{b}} \phi_{\underline{a}}^{\underline{b}} - \gamma_{\underline{a}} \equiv 0, \quad \text{mod } \mathcal{I}_{\text{tot}}.$$

As a result, by translating $r_{\underline{a}}$ to zero, the 1-forms $\gamma_{\underline{a}}$ are reduced, i.e.,

$$(3.8) \quad \gamma_{\underline{a}} = q_{ai} \omega^i + q_{ab} \theta^b.$$

After these reductions, the proposition below can be proved.

Proposition 3.1. *A causal structure with vanishing WSF curvature satisfying $f_{abc} = \lambda F_{abc}$ has vanishing SSF curvature.*

Proof. As was discussed, the condition $f_{abc} = \lambda F_{abc}$ results in the reduction of the 1-forms π_i according to equations (3.2), (3.3) and (3.5). In the case $p_{nn} \neq 0$, equation (3.6) it can be shown that ϕ_n can be reduced to (3.7) by normalizing p_{nn} to a non-zero value c . Consequently, the 1-forms γ_a can be reduced to (3.8).

Now that all the 1-forms π_i , ϕ_n and γ^a are reduced equation (3.4) can be written as

$$\begin{aligned} d\theta_a = & (\phi_a^b - \phi_0 \delta_a^b) \wedge \theta_b + c \omega^a \wedge \omega^n + (p_{ab} - p_{n0} \varepsilon_{ab}) \omega^0 \wedge \omega^b + p_{an} \omega^0 \wedge \omega^n \\ & - (p_{ab} + r_0 \varepsilon_{ab}) \theta^b \wedge \omega^0 - p_{nb} \varepsilon_{ac} \omega^b \wedge \omega^c - r_c \varepsilon_{ab} \theta^b \wedge \omega^c - r_n \theta^a \wedge \omega^n \end{aligned}$$

where c denotes the normalized value of p_{nn} . It is noted that the translations $r_{\underline{a}}, p_{n\underline{a}}, \varepsilon^{ab} p_{ab}$ to zero are used in the above equation. Similarly, express $d\omega^a$ and $d\omega^n$ by replacing γ^a and ϕ_n with their reduced forms in (2.37).

The reduced structure equations can be used to express the identity $d^2\theta_a = 0$. Collecting the coefficients of the 3-form $\theta^a \wedge \omega^b \wedge \omega^n$, it follows that

$$(3.9) \quad cF_{abc} + (r_{c;n} - r_{n;c} + \frac{1}{2}p_{cn} - p_{nc} - r_n q_{cn} - r_c r_n) \varepsilon_{ab} + p_{nb} \varepsilon_{ac} + p_{an} \varepsilon_{bc} + W_{abcn} = 0,$$

where $W_{abcn} = -W_{bacn}$ is a torsion term for Φ_{ab} as defined in (A.3) and (A.11). According to (A.20), the value of W_{abcn} is zero before the reduction. After reducing π_a and γ_a , it is found that

$$W_{abcn} = p_{bn} \varepsilon_{ac} - p_{an} \varepsilon_{bc}.$$

If $c \neq 0$, then equation (3.9) and the fact that F_{abc} is totally symmetric and trace-free imply $F_{abc} = 0$. This is a contradiction with the assumptions $f_{abc} = \lambda F_{abc}$ and F_{abc} being non-vanishing. As a result, c has to vanish. \square

Remark 3.2. The proposition above has an analogue in Finsler geometry. Recall that a Finsler metric is called Landsberg if the derivative of its Cartan torsion along geodesics is zero, i.e.,

$$\dot{I}_{ijk} = 0.$$

It can be shown that if the flag curvature of a Landsberg metric with non-vanishing Cartan torsion is constant, then it has to be zero [BCS00, Section 12.1]. By the proposition above, the condition $F_{abc;n} = \lambda F_{abc}$ can be used to translate $F_{abc;n}$ to zero. As a result of this translation, if the causal structure has vanishing WSF curvature, then its SSF curvature has to be zero. This observation suggests that causal structures satisfying $f_{abc} = \lambda F_{abc}$ are causal analogues of Landsberg spaces.

3.2. A rigidity theorem. In this section, under some conditions, a theorem involving the global vanishing of the Fubini form on a compact manifold is obtained.

As was discussed in Section 2.5.2, picking a representative of the conformal class of the quadratic form (2.35) results in the reduction (2.48). Consequently, translating s_i to zero, using the relation (2.49), results in further reductions (2.50).

The change in the structure equations after the reductions above can be obtained from (A.17) and (A.3) where the 1-form ϕ_n is replaced by its reduced form $-\phi_0 + s_{\underline{a}}\theta^a$ and π_i 's are replaced by their reduced form (2.50).

Assuming that the causal structure has vanishing WSF curvature, i.e., $W_{anab} = 0$, the Bianchi identity (2.46) implies that $W_{anbc} = 0$. Using $d^2\theta_a = 0$, it is found that p_{nn} is defined up to a scale as the infinitesimal action of the reduced structure group on it is given by

$$(3.10) \quad dp_{nn} + 4p_{nn}\phi_0 \equiv 0 \pmod{\mathcal{I}_{\text{tot}}}.$$

Hence, if it is non-zero, depending on its signature, it can be normalized to -1 or 1 , resulting in the reduction

$$(3.11) \quad \phi_0 = t_i\omega^i + t_{\underline{a}}\theta^a.$$

Using the reduced structure equations, the identity $d\phi_0$, can be used to give the infinitesimal action of the reduced structure group on $t_{\underline{a}}$ as

$$dt_{\underline{a}} - t_{\underline{b}}\phi^b_a - \frac{1}{2}\gamma_a \equiv 0 \pmod{\mathcal{I}_{\text{tot}}}.$$

Therefore, by translating $t_{\underline{a}}$ to zero, the 1-forms γ_a are reduced to

$$\gamma_a = q_{ai}\omega^i + q_{a\underline{b}}\theta^b.$$

Remark 3.3. Note that as a result of the conditions $F \neq 0$ and $p_{nn} \neq 0$ and the subsequent reductions, the first structure equations take the form

$$(3.12a) \quad d\omega^0 = -\theta_a \wedge \omega^a + T^0,$$

$$(3.12b) \quad d\omega^a = -\phi^a_b \wedge \omega^b - \theta^a \wedge \omega^n + T^a,$$

$$(3.12c) \quad d\omega^n = T^n,$$

$$(3.12d) \quad d\theta^a = -\phi^a_b \wedge \theta^b + T^a.$$

where

$$T^\epsilon = \frac{1}{2}T^\epsilon_{ij}\omega^i \wedge \omega^j + T^\epsilon_{\underline{a}i}\theta^a \wedge \omega^i + \frac{1}{2}T^\epsilon_{\underline{a}\underline{b}}\theta^a \wedge \theta^b,$$

for some coefficients $T^\epsilon_{ij}, T^\epsilon_{\underline{a}i}, T^\epsilon_{\underline{a}\underline{b}}$, with ϵ representing i and \underline{a} . As was mentioned, these are obtained from (A.17) and (A.3) after replacing the connection forms by their reduced expressions.

The structure group is reduced to

$$(3.13) \quad \left\{ \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & \mathbf{A} & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & \mathbf{A} \end{pmatrix} \middle| \mathbf{A} \in \text{O}(p, q) \right\}$$

Therefore, the bilinear forms

$$(3.14) \quad g = 2\omega^0 \circ \omega^n - \varepsilon_{ab}\omega^a \circ \omega^b, \quad h = 2\omega^0 \circ \omega^n - \varepsilon_{ab}\omega^a \circ \omega^b + \varepsilon_{ab}\theta^a \circ \theta^b$$

are preserved by the reduced structure group.

In the case $p_{nn} < 0$, the normalization $p_{nn} = -1$ can be achieved which results in the reductions that were explained above. Using $d^2\theta_a = 0$, using the structure equations after the reductions, the vanishing of the coefficient of $\theta^a \wedge \omega^b \wedge \omega^n$ implies the equation

$$(3.15) \quad F_{abc;nn} - F_{abc} = 0.$$

If M is compact, then all the components of F_{abc} have to be bounded.

Recall from Section 2.3.5, that a vector field tangent to the characteristic curves is annihilated by $\omega^0, \omega^a, \theta^a$. Now, let $s : \mathcal{C} \rightarrow \mathcal{P}$, be a section of the structure bundle, inducing the coframe $(s^*\omega^i, s^*\theta^a)$. The corresponding frame is denoted by $\frac{\partial}{\partial(s^*\omega^i)}, \frac{\partial}{\partial(s^*\theta^a)}$. The vector field $\frac{\partial}{\partial(s^*\omega^n)}$ is a tangent vector to the characteristic curves.

Definition 3.4. A characteristic curve γ , of a causal structure is called complete if for each section $s : \mathcal{C} \rightarrow \mathcal{P}$, the parametrization of γ with respect to $\frac{\partial}{\partial(s^* \omega^n)}$ can be extended to \mathbb{R} .

Now the stage is set to state the following theorem.

Theorem 3.5. *Let (M, \mathcal{C}) be a causal structure with vanishing WSF curvature over a compact manifold with non-zero Fubini cubic form. If the characteristic curves of \mathcal{C} are complete, then the scalar flag curvature of (M, \mathcal{C}) is non-negative.*

Proof. Restrict a section of the structure bundle, $s : \mathcal{C} \rightarrow \mathcal{P}$, to a characteristic curve and take a representative of the conformal class of the quadratic form (2.35). After carrying out the reductions, equation (3.15) gives a set of decoupled ODEs of the form $f'' - f = 0$ for each component F_{abc} of the Fubini cubic form. Assuming the completeness of the characteristic curves, it follows that the any non-zero solution of the ODE has to be unbounded. Since the manifold is compact, the components F_{abc} in the reduced structure equations are bounded and, consequently, have to vanish. \square

The above theorem has the following immediate corollary.

Corollary 3.6. *Let (M, \mathcal{C}) be a compact causal structure with vanishing WSF curvature and complete characteristic curves. If it has negative SSF curvature, then it is a flat pseudo-conformal structure.*

3.3. Semi-integrable Lie contact structures. In this section it is shown how a causal structure with vanishing WSF curvature descends to a Lie contact structures on \mathcal{N} . The construction is analogous to Grossman's notion of semi-integrable Segré structures arising from torsion-free path geometries [Gro00].

Let $p \in \mathcal{C}$ be a generic point, that is a point admitting an open neighborhood $U \subset \mathcal{C}$ foliated by characteristic curves. As a result, the quotient map $\tau : U \rightarrow \mathcal{N}^{2n-1}$ defines the space of characteristic curves in U .

Recall from Section 2.3.5 that a characteristic vector field \mathbf{v} spans the unique degenerate direction of ω^0 and $d\omega^0$, at every point i.e.,

$$\mathbf{v} \lrcorner \omega^0 = 0, \quad \mathbf{v} \lrcorner d\omega^0 = 0.$$

It follows that $\mathcal{L}_{\mathbf{v}} \omega^0 = 0$. Therefore, there exists a 1-form $\tilde{\omega}^0 \in \Gamma(T^* \mathcal{N})$ of rank of $2n - 1$ such that $\tau^* \tilde{\omega}^0 = \omega^0$. As a result, the quasi-contact structure on \mathcal{C} defined by ω^0 induces a contact distribution on \mathcal{N} . By abuse of notation, ω^0 is used to denote $\tilde{\omega}^0$ as the distinction is clear from the context.

Because

$$\mathbf{v} \lrcorner \omega^a = \mathbf{v} \lrcorner \theta^a = 0,$$

the 1-forms $\omega^0, \dots, \theta^{n-1}$ introduce a coframe on \mathcal{N} , and the $\{e\}$ -structure \mathcal{F} fibers over \mathcal{N} . To understand the resulting structure on \mathcal{N} , consider the flat model. Recall from Section 2.2 that the flat causal geometry corresponds to the null cone bundle of a hyperquadric $Q_{p+1, q+1} \subset \mathbb{P}^{n+2}$, and its space of characteristic curves is the space of isotropic 2-planes denoted by $G_0(2, n+4)$. The resulting structure was first studied by Sato and Yamaguchi in [SY89] referred to as Lie contact structures. As a parabolic geometry, this geometry is modeled on (B_n, P_2) or (D_n, P_2) , for $n \geq 4$ or $(D_3, P_{1,2})$ (see Section 2.4.1).

In this section it is shown how a causal structure with vanishing WSF curvature descends to a Lie contact structures on \mathcal{N} . As was discussed in Section 2.3.13, no attempt is made to introduce a Cartan connection for the Lie contact structure from the derived $\{e\}$ -structure.

First, a definition of Lie contact structures in terms of a field of Segré cones of type $(2, \mathbf{N})$ is given. Let \mathbb{V} be a $2\mathbf{N}$ -dimensional vector space with a decomposition $\mathbb{V} \cong \mathbb{R}^2 \otimes \mathbb{R}^{\mathbf{N}}$. An element $p \in \mathbb{V}$ is said to be of rank one if and only if $p = a \otimes b$. A Segré cone of type $(2, \mathbf{N})$, denoted by $S(2, \mathbf{N})$, is defined as the set of rank one elements of this decomposition. In terms of local coordinates $(z_1^1, \dots, z_{\mathbf{N}}^1, z_1^2, \dots, z_{\mathbf{N}}^2)$ adapted to this decomposition, the Segré cone is given by the points satisfying the homogeneous quadratic equations

$$z_{\mu}^1 z_{\nu}^2 - z_{\mu}^2 z_{\nu}^1 = 0,$$

where $1 \leq \mu, \nu \leq \mathbf{N}$. As a result, a Segré cone of type $(2, \mathbf{N})$ has a double ruling. A $(\mathbf{N} - 1)$ -parameter ruling by 2-planes $E_v \cong \mathbb{R}^2 \otimes v$ where $v \in \mathbb{R}^{\mathbf{N}}$. Such 2-planes are called α -planes

[AG96]. Another ruling of $S(2, \mathbf{N})$ is given by the 1-parameter family of \mathbf{N} -planes $u \otimes \mathbb{R}^{\mathbf{N}}$ where $u \in \mathbb{R}^2$. Such \mathbf{N} -planes are called β -planes.¹⁹

The proposition below is used to show how a Lie contact structure is induced on \mathcal{N} .

Proposition 3.7. *Given a causal structure with adapted coframe (ω^i, θ_a) , the span of the bilinear forms*

$$(3.16) \quad \Upsilon^{ab} = \theta^a \circ \omega^b - \theta^b \circ \omega^a$$

restricted to $H = \text{Ker } \omega^0$ is invariant under the action of the structure group G_5 . Moreover, \mathcal{C} has vanishing WSF curvature, if and only if the $\text{Span}\{\Upsilon^{ab}\}$ is invariant along characteristic curves.

Proof. By the matrix form of the structure group G_5 in (2.34), it follows that the span of Υ^{ab} modulo $I\{\omega^0\}$ is invariant under the action of G_5 .

Let \mathbf{v} denote a characteristic vector field. To verify the second claim it suffices to note that (2.37) implies

$$\mathcal{L}_{\mathbf{v}} \Upsilon^{ab} = A^{ab}_{cd} \Upsilon^{cd} + (\mathbf{v} \lrcorner \omega^n)(W^b_{ncn} \omega^a - W^a_{ncn} \omega^b) \circ \omega^c,$$

where $A^{ab}_{cd} = \mathbf{v} \lrcorner (2\phi_0 \delta_c^a \delta_d^b + \delta_d^b \phi_c^a - \delta_c^a \phi_d^b)$. Thus, if the condition $W^a_{nb n} = 0$ is satisfied, i.e., causal structure has vanishing WSF curvature, then the $\text{span}\{\Upsilon^{ab}\}$ is invariant along the characteristic curves. \square

As a result of the proposition above, the set of tangent vectors $\mathbf{u} \in H$ satisfying $\Upsilon^{ab}(\mathbf{u}, \mathbf{u}) = 0$ is well-defined and gives a Segré cone of type $(2, n-1)$ in the contact distribution $H \subset T\mathcal{N}$.

Definition 3.8. A manifold $N^{2\mathbf{N}+1}$ with a contact distribution H is said to have a Lie contact structure of signature (p, q) , $p + q = \mathbf{N}$ if there is a smoothly varying family of Segré cones $S(2, \mathbf{N}) \subset H_x$ such that at every point $x \in N$, its β -planes are endowed with conformally invariant inner product of signature (p, q) and are Lagrangian with respect to the symplectic 2-form that is induced by the contact 1-form on the contact distribution.

A Lie contact structure is called *semi-integrable* if there exists a 1-parameter family of foliations by Legendrian submanifolds with the property that given a point $x \in N$ and a β -plane in $S(2, \mathbf{N}) \subset H_x$, there passes a unique member of the family through x , tangent to that β -plane.

There are other equivalent definitions of a Lie contact structure discussed in [Zad09].

Remark 3.9. Semi-integrability condition gives rise to a foliation of the bundle $\tilde{\rho} : \mathcal{E}_{\beta} \rightarrow \mathcal{N}$ where \mathcal{E}_{β} denotes the bundle of β -planes of the Segré cones. Because β -planes are Lagrangian subspaces of the contact distribution, \mathcal{E}_{β} is a sub-bundle of $\text{LG}(\mathcal{N})$ where $\rho : \text{LG}(\mathcal{N}) \rightarrow \mathcal{N}$ is the bundle of Lagrangian-Grassmannians of the contact distribution H . Since, the bundle map $\tilde{\rho}$ is the restriction of ρ to the sub-bundle \mathcal{E}_{β} , by abuse of notation, it is denoted by ρ as well.

The theorem below gives a characterization of causal structures with vanishing WSF curvature in terms of semi-integrable Lie contact structures induced on their space of characteristic curves.

Theorem 3.10. *Given a causal structure (\mathcal{C}, M) with vanishing WSF curvature of signature $(p+1, q+1)$ its space of characteristic curves admits a semi-integrable Lie contact structure of signature (p, q) . Conversely, any semi-integrable Lie contact structure of signature (p, q) on a space \mathcal{N} induces a causal structure of signature $(p+1, q+1)$ with vanishing WSF curvature on its space of ruling Legendrian submanifolds.*

Proof. To see how \mathcal{N} is endowed with a Lie contact structure, firstly note that by Proposition 3.7 the contact distribution carries a Segré structure of type $(2, n-1)$. By the definition of Υ^{ab} , it follows that the β -planes, i.e., the 1-parameter ruling of the Segré cone, are given by

$$(3.17) \quad \text{Ker} \{A\theta^a + B\omega^a \mid a = 1, \dots, n-1\},$$

¹⁹The relation between these α - and β -planes in the section and those defined in Section 4.1 has to do with the fact that the null cones of a metric of split signature in four dimensions is a Segré cone of type $(2, 2)$.

$[A : B] \in \mathbb{P}^1$. Moreover, the inner product $\varepsilon_{ab}\theta^a \circ \theta^b$ is of signature (p, q) and its conformal class is well-defined on each β -plane, i.e., it is conformally invariant under the action of G_5 modulo $\mathcal{I}_{\text{bas}} = I\{\omega^0, \dots, \omega^n\}$. Finally, to show β -planes are Lagrangian consider a β -plane E_{β_1} corresponding to $[A : B] \in \mathbb{P}^1$ in (3.17). The tangent vectors (v_1, \dots, v_{n-1}) where $v_a = B \frac{\partial}{\partial \theta^a} - A \frac{\partial}{\partial \omega^a}$ define a basis for E_{β_1} . Because $d\omega^0 \equiv \varepsilon_{ab}\theta^a \wedge \omega^b$, modulo $I\{\omega^0\}$, it follows that $d\omega^0(v_a, v_b) = 0$. Hence, E_{β_1} is Lagrangian with respect to the conformal symplectic form $d\omega^0$.

As for the semi-integrability of these Lie contact structures, the following lemma will be useful.

Lemma 3.11. *Given a causal structure with vanishing WSF curvature, there is a canonical bundle diffeomorphism $\varpi : \mathcal{C} \rightarrow \mathcal{E}_\beta$ between the fiber bundles $\tau : \mathcal{C} \rightarrow \mathcal{N}$ and $\rho : \mathcal{E}_\beta \rightarrow \mathcal{N}$.*

Proof. Let $\gamma_t = (x(t); y(t))$ be a characteristic curve of \mathcal{C} which corresponds to the point $[\gamma] \in \mathcal{N}$. Let \mathcal{D} be the completely integrable distribution of $T\mathcal{C}$ given by the vertical tangent spaces $T_y\mathcal{C}_x := \text{Ker } \mathcal{I}_{\text{bas}}$ with $\mathcal{D}_t := T_{y(t)}\mathcal{C}_{x(t)}$. By the discussion above, it follows that, at $t = t_0$, $\tau_*\mathcal{D}_{t_0} \subset H_{[\gamma]}$ is isotropic, $(n-1)$ -dimensional and lies in the zero locus of Υ^{ab} , i.e. $\tau_*\mathcal{D}_{t_0}$ is a β -plane. By defining

$$(3.18) \quad \varpi(\gamma_{t_0}) = \tau_*\mathcal{D}_{t_0},$$

one obtains the desired diffeomorphism. \square

Continuing the proof of Theorem 3.10, note that the fibers \mathcal{C}_x , as the integral manifolds of \mathcal{D} , give a foliation of \mathcal{E}_β via the diffeomorphism ϖ . Because of the transversality of the fibers \mathcal{C}_x and characteristic curves γ passing through x , it follows that the subsets $\tau(\mathcal{C}_x) \subset \mathcal{N}$ are immersed as $(n-1)$ -dimensional submanifolds of \mathcal{N} . These submanifolds are Legendrian because it was shown that their tangent spaces, $\tau_*\mathcal{D}$, are β -planes, i.e., Lagrangian subspaces of the contact distribution which rule the Segré cone.

The converse part of the theorem can be shown as follows. Let ω^0 be a contact form. Since the Segré cones of type $(2, n-1)$ admit a 1-parameter ruling by $(n-1)$ -planes, one can find 1-forms $\{\eta^a, \zeta^a | a = 1, \dots, n-1\}$ complementing ω^0 in $T^*\mathcal{N}$ such that the Segré cone can be written as the vanishing set of the bilinear forms

$$\Upsilon^{ab} = \eta^a \circ \zeta^b - \eta^b \circ \zeta^a.$$

As a result, the 1-parameter family of ruling $(n-1)$ -planes of the Segré cone can be expressed as $\text{Ker } \{t\eta^a + (1-t)\zeta^a | a = 1, \dots, n-1\}$ where $t \in (0, 1)$. Let $\rho : \mathcal{E}_\beta \rightarrow \mathcal{N}$ denote the bundle of β -planes of the Segré cone. A point $p \in \mathcal{E}_\beta$ represents a β -plane of the contact distribution H which is Lagrangian with respect to the conformal symplectic structure on the contact distribution. Thus, the tautological bundle $K \subset T\mathcal{E}_\beta$ can be defined with fibers

$$K_p = \{v \in T_p\mathcal{E}_\beta \mid \rho_*|_p(v) \in p \subset T_{\rho(p)}\mathcal{N}\}.$$

As a result, the subspaces $\rho_*(K_p)$ rule the Segré cone $S(2, n-1) \subset H_{\rho(p)} \subset T_{\rho(p)}\mathcal{N}$. The fibers of $\rho : \mathcal{E}_\beta \rightarrow \mathcal{N}$ are one dimensional and parametrize the ruling subspaces $\rho_*(K_p)$. Let \mathbf{v} be a vertical vector field of the fibration $\rho : \mathcal{E}_\beta \rightarrow \mathcal{N}$ satisfying $\rho_*(\mathbf{v}) = \frac{\partial}{\partial t}$, and by abuse of notation, let a multiple of $\rho^*(\omega^0)$ be denoted by ω^0 as well. Hence, $\omega^0 \subset T^*\mathcal{E}_\beta$ is a quasi-contact 1-form. Defining $\omega^a = \rho^*(t\eta^a + (1-t)\zeta^a)$, it follows that $\mathcal{L}_{\mathbf{v}}\omega^a = \rho^*(\eta^a - \zeta^a)$, and, consequently,

$$\omega^1 \wedge \dots \wedge \omega^{n-1} \wedge \mathcal{L}_{\mathbf{v}}\omega^1 \wedge \dots \wedge \mathcal{L}_{\mathbf{v}}\omega^{n-1} \neq 0 \quad \text{mod} \quad I\{\omega^0\}.$$

The reason that it is non-zero, modulo $I\{\omega^0\}$, is that the Segré cone is a subset of the contact distribution.

Hence, locally, the tautological bundle is given by $K = \text{Ker } \{\omega^0, \dots, \omega^{n-1}\}$. As a result, the 1-forms $\omega^0, \omega^a, \theta_a = \mathcal{L}_{\mathbf{v}}\omega^a$ which are semi-basic with respect to the fibration $\rho : \mathcal{E}_\beta \rightarrow \mathcal{N}$ and the 1-form ω^n , which is vertical, span $T_p^*\mathcal{E}_\beta$, and one has

$$d\omega^a \equiv -\theta^a \wedge \omega^n + T^a_{bc}\theta^a \wedge \theta^b \quad \text{mod} \quad I\{\omega^0, \dots, \omega^{n-1}\}.$$

Since the Lie contact structure is assumed to be semi-integrable, the lift of its associated 1-parameter family of Legendrian submanifolds foliate \mathcal{E}_β by quasi-Legendrian submanifolds. The tangent distribution to this foliation is a corank sub-bundle of the tautological subbundle K given by $\text{Ker } \{\omega^0, \dots, \omega^n\}$. As a result, ω^i 's form a Pfaffian system which

implies $T^a_{bc} = 0$. Moreover, knowing that $K = \text{Ker} \{\omega^0, \dots, \omega^{n-1}\}$ and that ω^0 is semi-basic, one arrives at

$$d\omega^0 \equiv -h_{ab}\theta^a \wedge \omega^b, \quad \text{mod } I\{\omega^0\}.$$

In the identity $d^2\omega^0 = 0$, the vanishing of the 3-form $\theta^a \wedge \theta^b \wedge \omega^n$ implies that $h_{ab} = h_{ba}$.

Let M be the leaf space of the quasi-Legendrian foliation of \mathcal{E}_β , via the quotient map $\pi : \mathcal{E}_\beta \rightarrow M$. Because the tangent planes of the leaves are given by $\text{Ker} \{\omega^0, \dots, \omega^n\}$, the 1-forms $\{\omega^0, \dots, \omega^n\}$ are semi-basic with respect to the fibration $\pi : \mathcal{E}_\beta \rightarrow M$, and the 1-forms θ^a are vertical. Moreover, the coframe on \mathcal{E}_β is adapted to the flag (2.11). Note that the tautological bundle K is isomorphic to the tautological bundle introduced in Section 2.3.2.

According to Section 2.3, the complete integrability of the Pfaffian system \mathcal{I}_{bas} and the relations

$$\begin{aligned} \omega^0 &\equiv -\theta_a \wedge \omega^a, \quad \text{mod } I\{\omega^0\}, \\ \omega^a &\equiv -h^{ab}\theta_b \wedge \omega^n, \quad \text{mod } I\{\omega^0, \dots, \omega^{n-1}\}, \end{aligned}$$

where $\theta_a := h_{ab}\theta^b$ suffice to carry out the coframe adaptations and obtain the $\{e\}$ -structure that characterizes a causal structure. Note that the distribution given by $\text{Ker} \{\omega^0, \dots, \omega^{n-1}\}$ characterizes a causal structure according to Section 2.4.2. The one dimensional fibers of $\rho : \mathcal{E}_\beta \rightarrow \mathcal{N}$ are mapped to the characteristic curves of the causal structure.

Finally, since the vanishing set of the quadratic forms Υ^{ab} defined in (3.16) is preserved, the WSF curvature of the obtained causal structure has to vanish. \square

4. FOUR DIMENSIONS

In this section a class of causal structures on four dimensional manifolds is introduced for which a notion of half-flatness can be defined in terms of the existence of integral surfaces whose tangent planes in each tangent space give an envelope construction of the null cone at that point. As a result, one obtains a generalization of half-flat indefinite pseudo-conformal structures²⁰ in four dimensions. It is shown that the symmetry group of a non-flat causal structure of split signature, which does not descend to a pseudo-conformal structure, is at most 8-dimensional. Finally, the unique causal structure with 8-dimensional symmetry algebra is introduced.

4.1. Causal structures with ruled fibers. Given a causal structure of signature (2, 2), suppose the second fundamental form is normalized to $[\varepsilon_{ab}] = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$. Since F_{abc} is totally symmetric and trace-free, it has two independent components F_{111} and F_{222} . The Fubini-Pick invariant, $F := F_{abc}F^{abc}$, is given by $F = 8(F^1_{11})^2 - 8(F^2_{22})^2 = 8F^+F^-$, where

$$(4.1) \quad F^\pm = \frac{1}{\sqrt{2}}(F^1_{11} \pm F^2_{22}).$$

It is known that the vanishing of F implies that the projective surface is ruled [Sas99, GH79]. In this section, it is assumed that $F^+ = 0$. The case $F^- = 0$ can be treated analogously.

Define the following quantities

$$K^\pm := K^{11} \pm K^{12} = -F^\pm_{;1} + F^\pm_{;2} \quad L^\pm := \frac{1}{\sqrt{2}}(L^1 \pm L^2) = \frac{1}{2\sqrt{2}}(-K^\pm_{;1} + K^\pm_{;2}).$$

Because ε_{ab} is indefinite, one should keep in mind that raising or lowering of the index 2 is accompanied by a negative sign multiplication, e.g., $\theta_2 = -\theta^2$, or $F^2_{11} = -F_{211}$. Define

$$(4.2) \quad \omega^\pm := \frac{1}{\sqrt{2}}(\omega^1 \pm \omega^2), \quad \theta^\pm := \frac{1}{\sqrt{2}}(\theta_1 \mp \theta_2), \quad \gamma^\pm := \frac{1}{\sqrt{2}}(\gamma^1 \pm \gamma^2), \quad \pi_\pm := \frac{1}{\sqrt{2}}(\pi_1 \pm \pi_2).$$

²⁰In split signature, the terms half-flatness and self-duality are used to refer to pseudo-conformal structures with a 3-parameter family of null surfaces.

The structure equations (2.37) can be expressed as

$$(4.3a) \quad d\omega^0 = -2\phi_0 \wedge \omega^0 - \theta^- \wedge \omega^+ - \theta^+ \wedge \omega^-,$$

$$(4.3b) \quad d\omega^+ = -\gamma^+ \wedge \omega^0 - (\phi^1_2 + \phi_0 + \phi_3) \wedge \omega^+ - \theta^+ \wedge \omega^3, \\ -K^+ \theta^- \wedge \omega^0 - F^+ \theta^- \wedge \omega^-,$$

$$(4.3c) \quad d\omega^- = -\gamma^- \wedge \omega^0 - (-\phi^1_2 + \phi_0 + \phi_3) \wedge \omega^- - \theta^- \wedge \omega^3, \\ -K^- \theta^+ \wedge \omega^0 - F^- \theta^+ \wedge \omega^+,$$

$$(4.3d) \quad d\omega^3 = -\gamma^- \wedge \omega^+ - \gamma^+ \wedge \omega^- - 2\phi_3 \wedge \omega^3 - L^+ \theta^- \wedge \omega^0 - L^- \theta^+ \wedge \omega^0$$

$$(4.3e) \quad d\theta^+ = -\pi^+ \wedge \omega^0 - \pi_3 \wedge \omega^+ - (\phi^1_2 + \phi_0 - \phi_3) \wedge \theta^+ \\ + W^+ \omega^- \wedge \omega^3 + V^+ \omega^+ \wedge \omega^- - f^+ \theta^+ \wedge \omega^+,$$

$$(4.3f) \quad d\theta^- = -\pi^- \wedge \omega^0 - \pi_3 \wedge \omega^- - (-\phi^1_2 + \phi_0 - \phi_3) \wedge \theta^- \\ + W^- \omega^+ \wedge \omega^3 + V^- \omega^+ \wedge \omega^- - f^- \theta^+ \wedge \omega^+.$$

where

$$F^\pm = F^\pm_{;n}, \quad W^\pm = W_{1313} \mp W_{1323}, \quad V^\pm = W_{1312} \mp W_{2312},$$

Recall that the symmetric trace-free WSF curvature, W_{a3b3} , has two independent entries $W_{1313} = W_{2323}$ and $W_{1323} = W_{2313}$.

According to the structure equations (4.3), if in addition to the vanishing of F^+ , the quantity W^+ vanishes, then the Pfaffian system \mathcal{T} generated by $\{\omega^0, \omega^+, \theta^+\}$ becomes Frobenius. In fact, if $W^+ = 0$, then Bianchi identity (A.25) for W_{0abc} implies that $V^+ = 0$. This observation motivates the following definition.

Definition 4.1. A four dimensional causal structure is called *half-flat*, if the torsion coefficients F^+ , W^+ , or F^- , W^- vanish.

Remark 4.2. The above definition only makes sense when the causal structure is of signature (2,2), since in the Lorentzian case, i.e., signature (3,1), the Fubini-Pick invariant is given by

$$F = 8(F_{111}^2 + F_{222}^2),$$

which only vanishes if $F_{111}, F_{222} = 0$, i.e., the causal structure is equivalent to a conformal Lorentzian structure.

This term half-flat is justified as follows. In [Pen76], Penrose characterized self-dual pseudo-conformal structures of split signature, i.e., conformal structures for which the anti-self-dual part of the Weyl curvature vanishes, by the existence of a 3 parameter family of 2-surfaces. To explain his characterization, recall that for such pseudo-conformal structures on M , if the metric is given by $g = \omega^0 \circ \omega^3 - \omega^+ \circ \omega^-$ then at $x \in M$, there is a ruling of the null cone $\widehat{\mathcal{C}}_x \subset T_x M$ by the null 2-planes over which one has

$$(4.4) \quad \omega^0 + \lambda \omega^- = 0, \quad \omega^+ + \lambda \omega^3 = 0,$$

for some $\lambda \in \mathbb{R}$. In fact, λ parametrizes such ruling 2-planes (see [AG96]). There is another ruling of $\widehat{\mathcal{C}}_x$, given by the null 2-planes over which

$$(4.5) \quad \omega^0 + \mu \omega^+ = 0, \quad \omega^- + \mu \omega^3 = 0,$$

which are parametrized by μ . Assuming that M is oriented and that the null tetrad $(\omega^0, \omega^+, \omega^-, \omega^3)$ induces a positive orientation for M , the set of null 2-planes parametrized by λ and μ is referred to as α -planes and β -planes, respectively²¹. A 2-surface in M is called an α -surface or β -surface if its tangent space at each point is an α -plane or β -plane, respectively. A pseudo-conformal structure is called *half-flat* if there exists a 3-parameter family of α -surfaces in M , or equivalently, there exists a 3-parameter family of 2-surfaces with the property that at every point M , any given α -plane is tangent to a unique 2-surface in the family. Such pseudo-conformal structures are also called right-conformally flat [Pen76], self-dual [LM07], α -semi-integrable [AG96], or half-conformally flat [Bes08]. If M contains a 3-parameter family of β -surfaces then it is called half-flat as well. Alternatively, the terms left-conformally flat, anti-self-dual or β -semi-integrable can be used as well.

²¹In some literature α and β planes are defined the opposite way.

Penrose's characterization of half-flat pseudo-conformal structures can be generalized to causal structures in the following way.

Proposition 4.3. *Given a half-flat causal structure, there exists a 1-parameter family of foliations of M by 2-surfaces whose tangent planes at each point $x \in M$ rule the null cone \mathcal{C}_x .*

Proof. As mentioned before, if $W^+ = 0$, then the Bianchi identity (A.25) for W_{anbc} implies that $V^+ = 0$. As a result, the Pfaffian system \mathcal{T} generated by $\{\omega^0, \omega^+, \theta^+\}$ becomes completely integrable. The integral manifolds of \mathcal{T} are three dimensional submanifolds of \mathcal{C} which project to 2-surfaces in M . This is due to the fact that the tangent space of the integral manifolds are spanned by $\frac{\partial}{\partial \omega^-}, \frac{\partial}{\partial \omega^3}, \frac{\partial}{\partial \theta^-}$, where the tangent vector $\frac{\partial}{\partial \theta^-}$ is vertical with respect to the fibration $\pi : \mathcal{C} \rightarrow M$.

Now, consider the foliation of \mathcal{C} given by the integral curves of the Pfaffian system $\mathcal{U} = \{\omega^0, \omega^+, \omega^-, \omega^0, \theta^+\}$. The tangent directions of these integral curves are asymptotic directions of the fibers \mathcal{C}_x given by $\theta^+ = 0$. The first step is to show that at each point $x \in M$ these curves give a ruling of \mathcal{C}_x . Fix a point $x \in M$ and an integral curve of $\theta^+ = 0$ on \mathcal{C}_x denoted by $\rho(t)$. Let $\dot{\rho}(t)$ denote the vector $\frac{\partial}{\partial \theta^-}$ tangent to $\rho(t)$. The structure equations (4.3) imply

$$\mathcal{L}_{\dot{\rho}(t)}\omega^0 \wedge \mathcal{L}_{\dot{\rho}(t)}\omega^+ = f(x; \rho(t))\omega^0 \wedge \omega^+,$$

for some function $f(x; y)$. As a result, defining $P = \text{Ker } \{\omega^0, \omega^+\}$, the 2-plane $\pi_*(P) \subset T_x M$ is well-defined and can be represented by the bi-vector $\pi_*\left(\frac{\partial}{\partial \omega^-} \wedge \frac{\partial}{\partial \omega^3}\right)$. Moreover, because $\pi_*(P) \subset \widehat{T}_{\rho(t)}\mathcal{C}_x$, for all values of t , it follows that the cone over the curve $\rho(t)$, i.e., $\widehat{\rho(t)} \subset T_x M$, is a subset of $\pi_*(P)$. Hence, the integral curves of $\theta^+ = 0$ on \mathcal{C}_x are 2-plane in $T_x M$, which give a ruling of \mathcal{C}_x .

Using the foliation of the fibers \mathcal{C}_x by the Pfaffian system \mathcal{U} , one obtains the locally defined quotient map $\varrho : \mathcal{C} \rightarrow \mathcal{R}$. Moreover, because the integral curves of \mathcal{U} are curves in the fibers of the bundle $\mathcal{C} \rightarrow M$, the quotient space \mathcal{R} is a fiber bundle over M , denoted by (4.6)

$$\pi_{\text{rul}} : \mathcal{R} \rightarrow M,$$

and the fibers $\mathcal{R}_x := \pi_{\text{rul}}^{-1}(x)$ are 1-dimensional. From the previous discussion it follows that \mathcal{R}_x can be identified with a generating curve of the ruled fibers \mathcal{C}_x .

By the definition of \mathcal{R} , the integral manifolds of \mathcal{T} foliate \mathcal{R} by 2-surfaces and are transversal to the fibers \mathcal{R}_x . Under the projection π_{rul} one obtains a 1-parameter family of foliations of M by 2-surfaces. Moreover, the tangent planes of the 1-parameter family of 2-surfaces passing through each point $x \in M$, are given by $\pi_*\left(\frac{\partial}{\partial \omega^-} \wedge \frac{\partial}{\partial \omega^3}\right) \subset T_x M$ which, by our previous discussion, are the ruling 2-planes of \mathcal{C}_x .

Similarly, if the pair F^- and W^- vanish, the 2-surfaces are the projection of the integral manifolds of $\{\omega^0, \omega^-, \theta^-\}$. \square

Remark 4.4. Note that at each point $(x; y) \in \mathcal{C}$ the integral manifolds of \mathcal{U} are null with respect to the quadratic form $g = \omega^0 \circ \omega^3 - \omega^+ \circ \omega^-$ defined in (2.35). When the Fubini form vanishes, $[g]$ defines a pseudo-conformal structure of signature (2, 2) on M and the obtained 2-surfaces in M become its α -surfaces. In other words, the conditions

$$F^+ = 0, \quad \text{and} \quad W^+ = 0,$$

are the causal analogue of self-duality.

Remark 4.5. It can be shown that the 3-dimensional space of integral 3-folds foliating \mathcal{C} is equipped with a path structure whose paths are the image of the generating curve of the ruled surfaces \mathcal{C}_x under the quotient map. Conversely, given a path geometry, its space of paths is equipped with a half-flat causal structure [Mak16].

Comparing this observation with that of Grossman, [Gro00], it can be shown that what he refers to as the torsion of a path structure coincides with the fiber invariants, F^+, K^+, L^+ of a half-flat causal structure whose vanishing results in a half-flat conformal structure of split signature.

Example 4.6 (Product of two Finsler surfaces). Similar to Example 2.6.1, given two Finsler surfaces (N, Σ) and $(\tilde{N}, \tilde{\Sigma})$ one can consider the induced causal structure of split signature on their product space, denoted by $M = N \times_f \tilde{N}$, as defined in (2.60), where f is defined

over $N \times \tilde{N}$. According to structure equations (2.59), Finsler surfaces (N, Σ) and $(\tilde{N}, \tilde{\Sigma})$ are equipped with a unique coframing $(\eta^0, \eta^1, \zeta^1)$ and $(\tilde{\eta}^0, \tilde{\eta}^1, \tilde{\zeta}^1)$, and their associated scalars (I, J, K) and $(\tilde{I}, \tilde{J}, \tilde{K})$.

The coframe adaptation is rather similar to Example 2.6.1. For simplicity, the pull-back of coframing $(\tilde{\eta}^0, \tilde{\eta}^1, \tilde{\zeta}^1)$ is denoted by $(\eta^3, \eta^2, \zeta^2)$. A choice of 2-adapted coframing is given by

$$(4.7) \quad \begin{aligned} \omega^0 &= \frac{1}{\sqrt{2}}(-\eta^0 - f\eta^3), & \omega^1 &= \eta^1, & \omega^2 &= f\eta^2, & \omega^3 &= \frac{1}{\sqrt{2}}(-\eta^0 - f\eta^3), \\ \theta_1 &= \frac{1}{\sqrt{2}}\zeta_1 - \frac{1}{\sqrt{2}}f_{,1}\eta^3, & \theta_2 &= -\frac{1}{\sqrt{2}}\zeta_2 - \frac{1}{\sqrt{2}}\frac{f_{,2}}{f}\eta^3 + \frac{1}{\sqrt{2}}(f_{,0} + \frac{f_{,3}}{f})\eta^2, \\ \phi_0 &= -\phi_3 = -\frac{1}{2}f_{,0}\eta^3, & \gamma_1 &= -\frac{1}{\sqrt{2}}\zeta_1 - \frac{1}{\sqrt{2}}f_{,1}\eta^3, \\ \gamma_2 &= \frac{1}{\sqrt{2}}\zeta_2 + \frac{1}{\sqrt{2}}\frac{f_{,2}}{f}\eta^3 + \frac{1}{\sqrt{2}}(f_{,0} - \frac{f_{,3}}{f})\eta^2, & \phi &= \frac{f_{,1}}{f}\omega^2, & \pi_3 &= 0. \end{aligned}$$

After carrying out higher coframe adaptations and imposing the transformations (4.2), it follows that

$$\begin{aligned} F^\pm &= \frac{1}{2\sqrt{2}}(I \mp \tilde{I}), \\ W^\pm &= \frac{1}{4} \left(K - \frac{1}{f^2}\tilde{K} + (\ln f)_{,00} + (\ln f)_{,11} + \frac{1}{f}((\ln f)_{,03} + (\ln f)_{,30} - (\ln f)_{,0}(\ln f)_{,3}) \right. \\ &\quad \left. + \frac{1}{f^2}((\ln f)_{,33} - (\ln f)_{,22} - ((\ln f)_{,3})^2 + ((\ln f)_{,2})^2) \right) \mp \frac{1}{2f}(\ln f)_{,12}, \end{aligned}$$

where $(\ln f)_{,ij} := \frac{\partial}{\partial \eta^i} \frac{\partial}{\partial \eta^j} \ln f$. For instance if f is constant, $I = \tilde{I}$, then the condition $f^2 K = \tilde{K}$ results in a half-flat causal structure which has vanishing WSF curvature.

As another example suppose that f is a function on N , i.e., $f_{,2} = f_{,3} = 0$, which is also known as the warped product of N and \tilde{N} . Then, the half-flatness condition implies conformal flatness and is equivalent to the set of equations

$$I = \tilde{I}, \quad K = \frac{1}{f^2}\tilde{K} - (\ln f)_{,00} - (\ln f)_{,11}.$$

Example 4.7 (Cayley's cubic scroll). The causal structure introduced in Example 2.6.2 has ruled fibers $F^+ = 0$. The expression for W^+ is quite long. However, it can be checked that if

$$\begin{aligned} c_{111} &= c_{111}(x^1, x^3), & c_{12} &= e_1(x^1)e_2(x^2)e_3(x^3), \\ c_2 &= c_2(x^2), & c_1 &= c_1(x^1, x^3), & c_0 &= c_0(x^1, x^2), \end{aligned}$$

for some functions e_1, e_2 and e_3 then one has a half-flat causal structure.

4.2. Submaximal causal structures of split signature. A causal structure of split signature on a four dimensional manifold that does not descend to a pseudo-conformal structure is called *submaximal* if it is not flat and has the maximum dimension of symmetry algebra among all non-flat causal structures with non-zero Fubini cubic form.

The purpose of this section is to show the following proposition, making use of Definition 2.18 of a locally isotrivially flat causal structure.

Proposition 4.8. *The submaximal causal structure of split signature that does not descend to a pseudo-conformal structure has 8 dimensional symmetry algebra and is locally equivalent to the locally isotrivially flat causal structure whose null cones are projectively equivalent to Cayley's cubic scroll.*

Proof. First, it is shown that the maximum dimension of the symmetry algebra for a causal structure of split signature with non-zero Fubini cubic form is 8. Using the structure equations (4.3) or the infinitesimal group actions (2.39a), one obtains

$$(4.8a) \quad dF^\pm + F^\pm(\pm 3\phi^1_2 + \phi_0 - \phi_3) \equiv 0,$$

$$(4.8b) \quad dW^\pm + W^\pm(\pm 2\phi^1_2 - 4\phi_3) \equiv 0,$$

If $F^- \neq 0$, it can be normalized to one, resulting in the reduction

$$(4.9) \quad \phi_3 = \phi_0 + 3\phi^1_2 + r_i\omega^i + r_\mu\theta^\mu$$

where the range of i in this section is $\{0, +, -, 3\}$ and μ takes value in the set $\{+, -\}$. The reduction of ϕ_3 gives rise to the reduction of π_i 's and γ^μ 's as follows. First, one obtains

the infinitesimal group actions

$$\begin{aligned} dr_3 + \pi_3 &\equiv 0, \\ dr_\mu + \frac{1}{2}\pi_\mu &\equiv 0, \quad \text{mod } \mathcal{I}_{\text{tot}} \\ dr_{\underline{\mu}} + \frac{1}{2}\gamma_\mu &\equiv 0, \end{aligned}$$

Consequently, r_3, r_μ and $r_{\underline{\mu}}$ can be translated to zero resulting in reductions

$$\gamma_\mu = q_{\mu i}\omega^i + q_{\mu \underline{\nu}}\theta^\nu, \quad \pi_3 = p_{3i}\omega^i + p_{3\underline{\mu}}\theta^\mu, \quad \pi_\mu = p_{\mu i}\omega^i + p_{\mu \underline{\nu}}\theta^\nu.$$

Finally, the infinitesimal action

$$dp_{-\pm} + \pi_0 \equiv 0 \quad \text{mod } \mathcal{I}_{\text{tot}},$$

allows the reduction $\pi_0 = p_{0i}\omega^i + p_{0\underline{\mu}}\theta^\mu$.

As a result of the reductions above the structure bundle is reduced to 8 dimensions. If $F^+ \neq 0$, due to the reduction (4.9), the infinitesimal action (4.8a) on F^+ is expressed as

$$F^+ - 6F^+\phi^1{}_2 \equiv 0, \quad \text{mod } \mathcal{I}_{\text{tot}}.$$

Thus, the desired causal structure with 8 dimensional symmetry algebra satisfies $F^+ = 0$. Moreover, if the causal structure has non-zero WSF curvature, the relations (4.8b) result in further reductions.

Similar computations can be used to show that reduced structure group acts on $r_{\mu i}$, $r_{\mu \underline{\nu}}$, p_{ij} , $p_{i\underline{\mu}}$ via scaling. Hence, the only non-zero local invariant of the submaximal causal structure is F^- whose structure equations, after carrying out the reductions explained above, are given by

$$\begin{aligned} d\omega^0 &= -2\phi_0 \wedge \omega^0 - \theta^+ \wedge \omega^- - \theta^- \wedge \omega^+, \\ d\omega^+ &= -(2\phi_0 + 4\phi^1{}_2) \wedge \omega^+ - \theta^+ \wedge \omega^3, \\ d\omega^- &= -(2\phi_0 + 2\phi^1{}_2) \wedge \omega^- - \theta^- \wedge \omega^3 - \theta^+ \wedge \omega^+, \\ d\omega^3 &= -(2\phi_0 + 6\phi^1{}_2) \wedge \omega^3, \\ d\theta^+ &= 2\phi^1{}_2 \wedge \theta^+, \\ d\theta^- &= 4\phi^1{}_2 \wedge \theta^- \end{aligned} \tag{4.10}$$

The structure equations above are realized by the locally isotrivially flat causal structure whose null cones are Cayley's cubic scroll, i.e. its local defining function in affine coordinates is

$$F(x; y) = y^1 y^2 + \frac{1}{3}(y^1)^3$$

The conditions $F^+ \neq 0$ and $F^- = 0$ are no different and result in an equivalent causal structure with solvable 8-dimensional symmetry algebra. \square

APPENDIX A. FULL STRUCTURE EQUATIONS

The Appendix contains the computations needed to associate an $\{e\}$ -structure to a causal geometry. At the end, the symmetries and Bianchi identities for the torsion elements are expressed.

A.1. Prolongation and an $\{e\}$ -structure. Following the first order structure equations derived in (2.37), the method of equivalence can be used to associate a preferred choice of coframing to a causal structure. The first step involves prolonging the first order structure equations.

A.1.1. Prolonged structure group. Using the pseudo-connection in (2.37), one finds that the Cartan characters²² are

$$c_1 = 2n, \quad c_2 = n - 1, \quad c_i = n - i, \quad \text{for } 3 \leq i \leq n - 1.$$

²²see [Gar89, Olv95] for an account.

With respect to the basis $\{\omega^i, \theta^a, \phi_0, \phi_b^a, \phi_n, \gamma_a, \pi_n, \pi_a\}$, the prolonged structure group, $G_1^{(1)}$, is

$$(A.1) \quad \begin{pmatrix} \mathbb{I}_{2n} & 0 \\ \mathbf{R} & \mathbb{I}_p \end{pmatrix} \quad \text{such that} \quad \mathbf{R} = \begin{pmatrix} \mathbf{d} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & \mathbf{d} & 0 & 0 \\ \mathbf{f} & 0 & 0 & 0 \\ \mathbf{F} & \mathbf{f} & 0 & \mathbf{d} \end{pmatrix},$$

where $p = \frac{1}{2}(n^2 + n + 4)$, $\mathbf{E} \in \mathbb{R}^{n-1}$, and $\mathbf{d}, \mathbf{f} \in \mathbb{R}$. The Maurer-Cartan forms of the prolonged group can be written as

$$(A.2) \quad \begin{pmatrix} 0 & 0 \\ \mathbf{M} & 0 \end{pmatrix} \quad \text{where} \quad \mathbf{M} = \begin{pmatrix} \pi_0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & \pi_0 & 0 & 0 \\ \sigma & 0 & 0 & 0 \\ \tau_a & \sigma & 0 & \pi_0 \end{pmatrix},$$

Since $\dim G_1^{(1)} = n + 1$, the relation

$$\dim G_1^{(1)} < \sum_{j=1}^{n-1} j c_j,$$

holds for all $n \geq 3$. Thus, equations (2.37) are not involutive, and because an $\{e\}$ -structure is not obtained, according to Cartan's method of equivalence, one needs to prolong the structure equations.

A.1.2. Part of the second order structure equations. To avoid taking too much space for computations and for later use, the following 2-forms are introduced

$$(A.3) \quad \begin{aligned} \Omega^0 &= d\omega^0 + 2\phi_0 \wedge \omega^0 + \theta_a \wedge \omega^a, \\ \Omega^a &= d\omega^a + \gamma^a \wedge \omega^0 + (\phi_b^a + (\phi_0 + \phi_n)\delta_b^a) \wedge \omega^b + \theta^a \wedge \omega^n, \\ \Omega^n &= d\omega^n + \gamma_a \wedge \omega^a + 2\phi_n \wedge \omega^n, \\ \Theta^a &= d\theta^a + \pi^a \wedge \omega^0 + \pi_n \wedge \omega^a + (\phi_b^a - (\phi_0 - \phi_n)\delta_b^a) \wedge \theta^b \\ \Phi_0 &= d\phi_0 + \pi_0 \wedge \omega^0 + \frac{1}{2}\pi_a \wedge \omega^a - \frac{1}{2}\gamma_a \wedge \theta^a \\ \Phi_b^a &= d\phi_b^a + \phi_c^a \wedge \phi_b^c + \theta^a \wedge \gamma_b - \theta_b \wedge \gamma^a - \omega^a \wedge \pi_b + \omega_b \wedge \pi^a, \\ \Phi_n &= d\phi_n + \frac{1}{2}\pi_a \wedge \omega^a + \pi_n \wedge \omega^n + \frac{1}{2}\gamma_a \wedge \theta^a, \\ \Gamma_a &= d\gamma_a - \omega_a \wedge \pi_0 - \omega^n \wedge \pi_a - (\phi_b^a + (\phi_0 - \phi_n)\delta_b^a) \wedge \gamma_b, \\ \Pi_n &= d\pi_n - \theta^a \wedge \pi_a - 2\phi_n \wedge \pi_n, \\ \Pi_a &= d\pi_a - \theta_a \wedge \pi_0 - (\phi_b^a + (\phi_0 + \phi_n)\delta_b^a) \wedge \pi_b - \gamma_a \wedge \pi_n. \end{aligned}$$

As a result, structure equations (2.37) can be written as

$$(A.4) \quad \begin{aligned} \Omega^0 &= 0, \\ \Omega^a &= -F^a_{bc} \theta^b \wedge \omega^c - K^a_b \theta^b \wedge \omega^0, \\ \Omega^n &= -L_a \theta^a \wedge \omega^0, \\ \Theta_a &= -f_{abc} \theta^b \wedge \omega^c + \frac{1}{2}W_{anbc} \omega^b \wedge \omega^c + W_{bncn} \omega^c \wedge \omega^n. \end{aligned}$$

From the Lie algebra of the prolonged group in (A.2), it follows that

$$\Pi_n = -\sigma \wedge \omega^0 + P_n, \quad \Pi_a = -\tau_a \wedge \omega^0 - \sigma \wedge \omega_a + P_a,$$

where the 2-forms P_n and P_a are generated by $\{\omega^i, \theta^a, \phi_0, \phi_b^a, \phi_n, \gamma_a, \pi_n, \pi_a\}$. Note that the terms involving the 1-form π_0 in (A.2), are included within the definition of Γ_a, Π_n and Π_a .

Differentiating (2.37a), gives

$$(A.5) \quad 2\Phi_0 \wedge \omega^0 + \Theta_a \wedge \omega^a = 0.$$

As a result of (2.37d) and the symmetries in (2.38), it follows that $\Theta_a \wedge \omega^a = 0$, and therefore, $\Phi_0 = \zeta \wedge \omega^0$ for some 1-form ζ . By replacing

$$\pi_0 \mapsto \pi_0 - \zeta,$$

one arrives at

$$(A.6) \quad \Phi_0 = 0.$$

Differentiating (2.37c), and using the infinitesimal group action (2.39c), it is obtained

$$(A.7a) \quad \begin{aligned} \Gamma_a = & \frac{1}{2} W_{0aij} \omega^i \wedge \omega^j + E_{0ab0} \theta^b \wedge \omega^0 + \frac{1}{2} C_{abc} \theta^b \wedge \theta^c \\ & - F_{abc} \gamma^b \wedge \theta^c - L_a \pi_n \wedge \omega^0 + \tau_{ab} \wedge \omega^b + \xi_a \wedge \omega^n, \end{aligned}$$

$$(A.7b) \quad \Phi_n = W_{nnij} \omega^i \wedge \omega^j - \frac{1}{2} \ell_a \theta^a \wedge \omega^0 + \nu \wedge \omega^n + \frac{1}{2} \xi_a \wedge \omega^a,$$

for some coefficients E_{nab0}, C_{abc} , where $\ell_a = L_{a,n}$, and the yet undetermined 1-forms ν, ξ_a , and $\tau_{ab} = \tau_{ba}$ which are congruent to zero modulo \mathcal{I}_{tot} . The reason for using W_{nnij} instead of $\frac{1}{2} W_{nnij}$ in (A.7b) becomes clear when the replacements (A.13) are imposed.

By differentiating (2.37b), it follows that

$$\xi_a = k_{ab} \theta^b,$$

where $k_{ab} := K_{ab;n}$.

Now Φ^a_b can be determined as a result of differentiating (2.37b) and replacing Φ_0 and Φ_n by the expressions (A.6) and (A.7b). Using the infinitesimal group action on K_{ab}, F_{abc} given by (2.39a) and (2.39b), and their symmetries in (2.38), and (2.37d), the derivative of (2.37b) yields

$$(A.8) \quad \begin{aligned} 0 \equiv & (\Phi^a_b + \Phi_n \delta^a_b) \wedge \omega^b + K_{ab} \theta^b \wedge \theta_c \wedge \omega^c - F_{abc} f^b_{de} \theta^d \wedge \omega^e \wedge \omega^c \\ & + F_{abc;d} \omega^d \wedge \theta^b \wedge \omega^c + F_{abc;\underline{d}} \theta^d \wedge \theta^b \wedge \omega^c + F_{abc} F^c_{de} \theta^b \wedge \theta^d \wedge \omega^e \\ & + \frac{1}{2} F_{abc} W^b_{nde} \omega^d \wedge \omega^e \wedge \omega^c, \quad \text{mod } I\{\omega^0, \omega^n\}. \end{aligned}$$

Multiplied by the $(n-1)$ -form $\eta = \theta^1 \wedge \dots \wedge \theta^{n-1}$, the above equation results in

$$(\Phi^a_b + \Phi_n \delta^a_b) \wedge \omega^b \wedge \eta \equiv \frac{1}{2} - F_{aeb} W^b_{ncb} \omega^c \wedge \omega^d \wedge \omega^b \wedge \eta, \quad \text{mod } I\{\omega^0, \omega^n\}.$$

By Cartan's lemma, it follows that

$$(\Phi^a_b + \Phi_n \delta^a_b) \wedge \eta \equiv -(\zeta^a_{bc} \wedge \omega^c + \frac{1}{2} F_{abe} W^e_{ncd} \omega^c \wedge \omega^d) \wedge \eta, \quad \text{mod } I\{\omega^0, \omega^n\},$$

for some 1-form ζ^a_{bc} , where $\zeta^a_{bc} \equiv \zeta^a_{cb}$ modulo \mathcal{I}_{tot} .

Fixing $c = C$, after taking the wedge product of $\Phi^a_b + \Phi_n \delta^a_b$ with all the 1-forms ω^i , such that $i \neq C$, and using (A.7b), the result is

$$\Phi^a_b \wedge \omega^0 \wedge \dots \wedge \widehat{\omega^C} \wedge \dots \wedge \omega^n \wedge \eta = \zeta^a_{bC} \wedge \omega^C \wedge \omega^0 \wedge \dots \wedge \widehat{\omega^C} \wedge \dots \wedge \omega^n \wedge \eta.$$

Carrying out the same procedure for Φ^b_a , using $\Phi_{ab} + \Phi_{ba} = 0$, it follows that

$$(\zeta_{abC} + \zeta_{baC}) \wedge \omega^C \wedge \omega^0 \wedge \dots \wedge \widehat{\omega^C} \wedge \dots \wedge \omega^n \wedge \eta = 0,$$

and therefore, $\zeta_{abc} + \zeta_{bac} \equiv 0$ modulo \mathcal{I}_{tot} . However, since $\zeta^a_{bc} \equiv \zeta^a_{cb}$, one obtains $\zeta^a_{bc} \equiv 0$ modulo \mathcal{I}_{tot} .

Consequently, equation

$$(A.9) \quad \Phi_{ab} = \frac{1}{2} W_{abij} \omega^i \wedge \omega^j + E_{abci} \theta^c \wedge \omega^i + \frac{1}{2} C_{abcd} \theta^c \wedge \theta^d + \lambda_{ab} \wedge \omega^0 + \kappa_{ab} \wedge \omega^n,$$

is obtained where $\lambda_{ab} = -\lambda_{ba}, \kappa_{ab} = -\kappa_{ba}$ and $\lambda_{ab}, \kappa_{ab} \equiv 0$ modulo \mathcal{I}_{tot} .

Differentiating (2.37b), and using (A.7), and (A.9), it is straightforward to find

$$(A.10) \quad \kappa_{ab} = 0, \quad \nu = 0, \quad \tau_{ab} = E_{0acb} \theta^c + K_{ab} \pi_n - F_{ab}{}^c \pi_c, \quad \lambda_{ab} = 0,$$

for some E_{0abc} .

Equations (A.10), (A.7), (A.9) and (A.6) can be combined to give

$$(A.11) \quad \begin{aligned} \Phi_0 &= 0, \\ \Phi_{ab} &= \frac{1}{2} W_{abij} \omega^i \wedge \omega^j + E_{abci} \theta^c \wedge \omega^i + \frac{1}{2} C_{abcd} \theta^c \wedge \theta^d \\ \Phi_n &= W_{nnij} \omega^i \wedge \omega^j - \frac{1}{2} \ell_a \theta^a \wedge \omega^0 + \frac{1}{2} k_{ab} \theta^b \wedge \omega^a, \\ \Gamma_a &= \frac{1}{2} W_{0aij} \omega^i \wedge \omega^j + E_{0abi} \theta^b \wedge \omega^i + \frac{1}{2} C_{abc} \theta^b \wedge \theta^c \\ &\quad - F_{abc} \gamma^b \wedge \theta^c - L_a \pi_n \wedge \omega^0 + K_{ab} \pi_n \wedge \omega^b - F_{ab}{}^c \pi_c \wedge \omega^b. \end{aligned}$$

The quantities W_{ijkl} , E_{ijkl} , C_{abcd} and C_{abc} are not expressed until the replacement (A.13) is made which makes the computation slightly easier.

A.1.3. Reduction of the prolonged group. The infinitesimal action of $G_1^{(1)}$ on W_{nn0n} and W_{nn0a} is given by

$$dW_{nn0n} + \frac{1}{2}\sigma \equiv 0 \quad dW_{nn0a} + \frac{1}{4}\tau_a \equiv 0$$

modulo the ideal generated by the 1-forms $\{\omega^i, \theta_a, \phi_0, \phi_n, \phi^a_b, \gamma_a, \pi_n, \pi_a\}_{i=0, a=1}^{n, n-1}$.

Consequently, the normalizations $W_{nn0n} = 0$, $W_{nn0a} = 0$, reduce $G_1^{(1)}$ to the group $G_2^{(1)}$ containing the one parameter \mathbf{d} in (A.1), which corresponds to the 1-form π_0 . As a result, the expression for Φ_n changes to

$$(A.12) \quad \Phi_n = W_{nnab} \omega^a \wedge \omega^b + 2W_{nnan} \omega^a \wedge \omega^n - \frac{1}{2}\ell_a \theta^a \wedge \omega^0 + \frac{1}{2}k_{ab} \theta^b \wedge \omega^a$$

One sees that the action of the parameter \mathbf{d} on the torsion coefficients in the 2-forms $\Theta_0, \dots, \Phi_0, \dots, \Gamma_{n-1}$ a priori cannot be used to reduce the prolonged structure group to the trivial group, unless certain non-vanishing conditions are assumed.

The only part left, is to see whether any reduction is possible via the torsion coefficients in Π_n, \dots, Π_0 .

A.1.4. An $\{e\}$ -structure. In order to make computing Π_0, \dots, Π_n easier the replacements

$$(A.13) \quad \pi_0 \mapsto \pi_0 + \frac{1}{2}\ell_a \theta^a, \quad \pi_a \mapsto \pi_a - \frac{1}{2}k_{ab} \theta^b + W_{nnab} \omega^b, \quad \pi_n \mapsto \pi_n - 2W_{nnan} \omega^a.$$

are imposed which result in

$$\Phi_0 + \Phi_n = 0.$$

Similarly to Section A.1.2, by differentiating the 2-forms Γ_a, Φ^a_b in (A.11), Φ_n in (A.12) and $\Phi_0 = -\Phi_n$, and carrying out tedious but straightforward calculations, it follows that

$$(A.14) \quad \begin{aligned} \Pi_n &= \frac{1}{2}W_{nij} \omega^i \wedge \omega^j + E_{nai} \theta^a \wedge \omega^i + \sigma_{ni} \wedge \omega^i, \\ \Pi_a &= \frac{1}{2}W_{aij} \omega^i \wedge \omega^j + E_{abi} \theta^b \wedge \omega^i - F_{ab}{}^c \pi_c \wedge \theta^b + \sigma_{ai} \wedge \omega^i, \end{aligned}$$

where $\sigma_{ab} = \sigma_{ba}$, $\sigma_{na} = \sigma_{an}$ are congruent to zero modulo \mathcal{I}_{tot} .

The expressions for the torsion coefficients are listed in (A.20). It turns out that without assuming certain non-vanishing conditions on the torsion coefficients, the infinitesimal action of the parameter \mathbf{d} on the torsion terms does not reduce the prolonged group.

Since, the prolongation of the reduced group $G_2^{(1)}$ is trivial, it follows that the set of 1-forms

$$\{\omega^i, \theta^a, \phi_0, \phi^a_b, \phi_n, \gamma_a, \pi_i\}_{i=0, a=1}^{n, n-1}$$

constitutes an $\{e\}$ -structure for \mathcal{C} .

A.1.5. The third order structure equations. Similarly to (A.3), defining the 2-form

$$(A.15) \quad \Pi_0 = d\pi_0 - 2\phi_0 \wedge \pi_0 - \gamma^a \wedge \pi_a,$$

helps making the expressions more compact.

It is a matter of computation, similar to Section A.1.4, to find

$$(A.16) \quad \Pi_0 = \frac{1}{2}W_{0ij} \omega^i \wedge \omega^j + E_{0ai} \theta^a \wedge \omega^i - K_{ab} \pi^a \wedge \theta^b - L_a \pi_n \wedge \theta^a + \sigma_{0i} \wedge \omega^i,$$

where 1-forms σ_{ij} defined in (A.14) satisfy $\sigma_{ij} = \sigma_{ji}$. Note that the symmetry of σ_{ij} is a result of imposing $\Phi_0 + \Phi_n = 0$. Expressions for σ_{ij} are listed in (A.22).

A.1.6. *Full structure equations.* The full structure equations after replacement (A.13) using the 2-forms (A.3) are expressed as follows.

$$\begin{aligned}
\Omega^0 &= 0, \\
\Omega^a &= -F^a_{bc} \theta^b \wedge \omega^c - K^a_b \theta^b \wedge \omega^0, \\
\Omega^n &= -L_a \theta^a \wedge \omega^0, \\
\Theta_a &= -f_{abc} \theta^b \wedge \omega^c - \frac{1}{2} k_{ab} \theta^b \wedge \omega^0 - W_{nnab} \omega^0 \wedge \omega^b \\
&\quad + \frac{1}{2} W_{anbc} \omega^b \wedge \omega^c + W_{bnca} \omega^c \wedge \omega^n, \\
\Phi_0 &= -\frac{1}{2} W_{nnab} \omega^a \wedge \omega^b + \frac{1}{2} \ell_a \theta^a \wedge \omega^0 - \frac{1}{4} k_{ab} \theta^b \wedge \omega^a, \\
\Phi_{ab} &= \frac{1}{2} W_{abij} \omega^i \wedge \omega^j + E_{abci} \theta^c \wedge \omega^i + \frac{1}{2} C_{abcd} \theta^c \wedge \theta^d \\
\Phi_n &= \frac{1}{2} W_{nnab} \omega^a \wedge \omega^b - \frac{1}{2} \ell_a \theta^a \wedge \omega^0 + \frac{1}{4} k_{ab} \theta^b \wedge \omega^a, \\
\Gamma_a &= \frac{1}{2} W_{0aij} \omega^i \wedge \omega^j + E_{0abi} \theta^b \wedge \omega^i + \frac{1}{2} C_{abc} \theta^b \wedge \theta^c \\
&\quad - F_{abc} \gamma^b \wedge \theta^c - L_a \pi_n \wedge \omega^0 + K_{ab} \pi_n \wedge \omega^b - F_{ab}^c \pi_c \wedge \omega^b, \\
\Pi_n &= \frac{1}{2} W_{nij} \omega^i \wedge \omega^j + E_{nai} \theta^a \wedge \omega^i + \sigma_{ni} \wedge \omega^i, \\
\Pi_a &= \frac{1}{2} W_{aij} \omega^i \wedge \omega^j + E_{abi} \theta^b \wedge \omega^i - F_{ab}^c \pi_c \wedge \theta^b + \sigma_{ai} \wedge \omega^i, \\
\Pi_0 &= \frac{1}{2} W_{0ij} \omega^i \wedge \omega^j + E_{0ai} \theta^a \wedge \omega^i - K_{ab} \pi^a \wedge \theta^b - L_a \pi_n \wedge \theta^a + \sigma_{0i} \wedge \omega^i,
\end{aligned} \tag{A.17}$$

The expressions of the quantities on the right hand side are given in (A.20), (A.22) and (A.23). When differentiation (2.37b), the vanishing of 3-forms $\theta^b \wedge \theta^c \wedge \omega^d$, $\theta^b \wedge \theta^c \wedge \omega^0$ combined with the expression of C_{abcd} , and C_{abc} in (A.20), yield the relations

$$(A.18a) \quad F_{abc, \underline{d}} - F_{abd, \underline{c}} = \frac{1}{2} (K_{ac} \varepsilon_{bd} + K_{bc} \varepsilon_{ad} - K_{ad} \varepsilon_{bc} - K_{bd} \varepsilon_{ac}),$$

$$(A.18b) \quad K_{ab, \underline{c}} - K_{ac, \underline{b}} = 2(\varepsilon_{ab} L_c - \varepsilon_{ac} L_b) + F_{abd} K^d_c - F_{acd} K^d_b,$$

Moreover, the vanishing of the 3-forms $\theta_a \wedge \theta_b \wedge \omega^0$, in the exterior derivative of (2.37c) implies $L_{a; \underline{b}} = L_{b; \underline{a}}$. This equation together with the contraction of Equations (A.18) give the following Bianchi identities among the fiber invariants of \mathcal{C}

$$(A.19a) \quad K_{ab} = \frac{2}{n-1} F^c_{ab; \underline{c}},$$

$$(A.19b) \quad L_a = \frac{1}{2(n-2)} (F_{abc} K^{bc} - K_a^b{}_{; \underline{b}}),$$

$$(A.19c) \quad L_{a; \underline{b}} = L_{b; \underline{a}}.$$

A.2. **Quantities in (A.17).** The expression for the torsion coefficients are as follows:

$$\begin{aligned}
E_{abc0} &= \frac{1}{2} (K_{bc; a} - K_{ac; b} - K_{ad} f^d_{bc} + K_{bd} f^d_{ac}), \quad E_{abcn} = 0 \\
E_{abcd} &= F_{bcd; a} - F_{acd; b} + F_{bde} f^e_{ca} - F_{ade} f^e_{cb} \\
C_{abcd} &= F_{bce} F^e_{ad} - F_{ace} F^e_{bd} + \frac{1}{2} (K_{bc} \varepsilon_{ad} + K_{ad} \varepsilon_{bc} - K_{ac} \varepsilon_{bd} - K_{bd} \varepsilon_{ac}) \\
E_{0ab0} &= -L_{b; a} - L_d f^d_{ab}, \quad E_{0abn} = \frac{1}{2} k_{ab}, \quad C_{abc} = L_c \varepsilon_{ab} - L_b \varepsilon_{ac} \\
E_{0abc} &= \frac{1}{2} (K_{ab; c} + K_{bc; a} + K_{ad} f^d_{bc} + K_{cd} f^d_{ab}) - F_{abc; 0} - \frac{1}{2} k_{bd} F^d_{ac}
\end{aligned} \tag{A.20}$$

The quantities $W_{ijkl} = -W_{ijlk}$ satisfy the following identities:

$$\begin{aligned}
W_{an0b} &= -W_{nnab}, \quad W_{an0n} = 0, \quad W_{anbn} = W_{bnan}, \quad W^a_{nan} = 0, \quad W_{[a|n|bc]} = 0, \\
W_{abcn} &= W_{cnab} + F_{acd} W^d_{nbn} - F_{bcd} W^d_{nan}, \quad W_{a[bcd]} + F_{ae[b|} W_{en|cd]} = 0, \\
W_{abcd} - W_{cdab} &= F_{aec} W^e_{nbd} - F_{aed} W^e_{bnc} - F_{bec} W^e_{nad} + F_{bed} W^e_{nac} \\
W_{ab0c} &= -W_{0cab} + F_{acd} W^d_{n0b} - F_{bcd} W^d_{n0a} + \frac{1}{2} K_{cd} W^d_{nba} \\
&\quad + \frac{1}{2} (K_{bd} W^d_{nca} - K_{ad} W^d_{ncb}) \\
W_{0abn} &= -W_{nnab} - K_{(a|d} W^d_{n|b)n}, \quad W_{ab0n} = -2W_{nnab} + K_{[a|d} W^d_{n|b]n} \\
W_{0[abc]} &= 0, \quad W_{0a0n} = L_c W^c_{nab}, \quad W_{0[a|0|b]} = \frac{1}{2} L_c W^c_{nab}, \quad W_0{}^a{}_{0a} = 0.
\end{aligned} \tag{A.21}$$

According to the symmetries above, the independent components of W_{ijkl} are W_{abcd} , W_{anbc} , W_{0abc} , W_{nabn} , W_{0a0b} and W_{nnab} . It is shown in the A.3 that W_{ijkl} is generated by W_{anbn} .

The 1-forms $\sigma_{ij} = \sigma_{ji}$ are

$$\begin{aligned}
 \sigma_{nn} &= 0, \\
 \sigma_{an} &= W_{anbn}\gamma^b, \\
 \sigma_{a0} &= 3W_{nnab}\gamma^b - \ell_a\pi_n - \frac{1}{2}k_{ab}\pi^b, \\
 \sigma_{0n} &= 0, \\
 \sigma_{ab} &= -W_{(a|n|b)c}\gamma^c + \frac{1}{2}k_{ab}\pi_n - f_{abc}\pi^c, \\
 \sigma_{00} &= \frac{1}{n-2} \left(2W_{0bba} + K^{cd}W_{dnca} - L^c W_{cnan} \right) \gamma^a \\
 &\quad - \frac{2}{n-1} L^a{}_{;a} \pi_n - \frac{1}{n-1} \left(K^{ac}{}_{;a} + K_{ab} f^{abc} \right) \pi_c.
 \end{aligned}
 \tag{A.22}$$

The quantities $E_{iaj} = E_{jai}$ are given by

$$\begin{aligned}
 E_{nan} &= \frac{2}{n-1} \varepsilon^{bc} W_{0bca}, \\
 E_{nab} &= 3W_{nnab} + \frac{1}{2}k_{ab;n} + K_{(a|d} W^d{}_{n|b)n}, \\
 E_{0ab} &= -W_{0a0b} + 2W_{nnbd} K^d{}_a - \ell_{a;b} - \ell_d f^d{}_{ab} - \frac{1}{2}k_{ab;0} - \frac{1}{4}k_a{}^c k_{bc}, \\
 E_{abc} &= W_{(a|b0|a)} - W_{(a|n|c)d} K_{db} - W_{ancn} L_b + \frac{1}{2}K_{(a|b;n|c)} - F_{abc;n0} \\
 &\quad - \frac{1}{2}F_{acd;n} K_{db;n} - \frac{1}{2}K_{(a|d;n} F_{db|c);n} + \left(\frac{1}{2}L_c W_{cnbn} + \frac{1}{2}\ell_{b;n} \right) \varepsilon_{ac} \\
 E_{0an} &= -\frac{1}{2}L_c W_{cnan} - \frac{1}{2}\ell_{a;n}, \\
 E_{0b0} &= \frac{1}{n-1} \left(-2F_{abc} W_{0d0c} + W_{0aad} K_{db} - K^{ad} W_{nand} L_b + L_a E_{nba} - L_{a;d} f^{ad}{}_b \right. \\
 &\quad \left. - (f^a{}_{bc} L^c - \varepsilon^{ca} L_{b;c})_{;a} L^c f_{acd} f^{ad}{}_b - \frac{1}{2} (K^a{}_{d;a} + K^{ae} f_{ade}) k^d{}_b \right. \\
 &\quad \left. - (K^a{}_{b;a} + K^{ad} f_{adb})_{;0} + L^d W_{nndb} \right)
 \end{aligned}
 \tag{A.23}$$

The quantities W_{ijk} satisfy

$$W_{ijk} + W_{kij} + W_{jki} = 0, \quad W_{ijk} = -W_{ikj}$$

and are obtained from W_{ijkl}, F_{abc} as follows

$$\begin{aligned}
 W_{nan} &= -\frac{1}{n-2} (W_{anba;n} - W_{anbn;a} - F_{adb;n} W_{dnan}), \\
 W_{nab} &= 2W_{nnab} + k_{[a|c} W_{cn|b]n} \\
 W_{n0n} &= \frac{1}{(n-2)(n-1)} (W_{cdcd;n} - E_{cdec} W_{dnen} - E_{cded} W_{encn} - W_{adan;d} + W_{addn;a}) \\
 &\quad + \frac{1}{2(n-1)} k_{cd} W_{ancn}, \\
 W_{abn} &= \varepsilon_{ab} W_{n0n} + W_{anbn;0} - W_{nnab;n} - \frac{1}{2}k_{ac} W_{cnbn} \\
 W_{a0n} &= \frac{1}{n-2} \left(-W_{nnba;b} - W_{bnba;0} + \frac{1}{2}k_{ed} W_{dnea} + W_{0dda;n} - W_{0ddn;a} \right. \\
 &\quad \left. + W_{0dan;d} - E_{eda} W_{dnen} + E_{ede} W_{dnan} + E_{0edn} W_{dnea} \right. \\
 &\quad \left. - K_{ad} W_{ndn} + F_{ade} W_{den} \right) \\
 W_{n0a} &= \frac{1}{2} W_{a0n} - \frac{1}{2} \ell_b W_{bnan}, \\
 W_{abc} &= \varepsilon_{ab} W_{n0c} - \varepsilon_{ac} W_{n0b} - W_{nnab;c} + W_{nnac;b} + W_{anbc;0} \\
 &\quad - \frac{1}{2} K_{ad} W_{dnbc} + f_{acd} W_{nndb} - f_{abd} W_{nn dc}, \\
 W_{0ab} &= -2W_{nnab;0} + \ell_c W_{cnab} - k_{[a|d} W_{nn c|b]}, \\
 W_{00n} &= \frac{1}{(n-1)(n-2)} \left(W_{ab0[b;a]} - \frac{1}{2} W_{abab;0} + \frac{1}{2} E_{abe0} W_{enba} + \frac{1}{2} E_{abe0} W_{nnea} \right) \\
 &\quad + \frac{1}{n-1} (E_{0adn} W_{nnda} - E_{0ad0} W_{dnad} + L_a W_{nan}), \\
 W_{a0b} &= \varepsilon_{ab} W_{00n} + W_{0a0b;n} + W_{0abn;0} - E_{0adn} W_{nndb}, \\
 &\quad + E_{0ad0} W_{dnbn} - L_a W_{nbn} - K_{ab} W_{n0n} + F_{abc} W_{c0n}, \\
 W_{00b} &= -W_{0aab;0} + W_{0a0b;a} + E_{0ab0} W_{dba} + E_{0ada} W_{d0b} \\
 &\quad - E_{0adb} W_{d0a} - L_a W_{nba} - K_{ab} W_{n0a} + F_{abd} W_{d0a}
 \end{aligned}
 \tag{A.24}$$

In the above expressions the indices are lowered using ε^{ab} and the summation is taken over repeated indices.

A.3. Bianchi identities for W_{ijkl} . The following relations hold among the torsion elements W_{ijkl} in (A.17).

$$\begin{aligned}
 W_{anbc} &= \frac{2}{3} \left(W_{an[b|n;|\underline{c}]} + F_{a[c}{}^d W_{b]ndn} \right) \\
 &\quad + \frac{4}{3(n+2)} \left(\varepsilon_{a[c]} W_{n|b]n;\underline{d}}^d + \varepsilon_{a[b} F^{de}{}_{c]} W_{dnen} \right) \\
 W_{nnab} &= \frac{1}{2(n+3)} \left(-W_{nab;\underline{d}}^d + F^{cd}{}_a W_{cndb} - F^{cd}{}_b W_{cnda} \right), \\
 W_{abcd} &= W_{[a|ncd;|\underline{b}]} - f^e{}_{ac} f_{bde} + f^e{}_{ad} f_{bce} + W_{ane[c} F_{b]d}{}^e - W_{dne[c} F_{b]a}{}^e \\
 &\quad - \varepsilon_{a[c]} \left(W_{nn|d]b} - \frac{1}{2} k_{[d]b;n} + W_{d]nen} K_b{}^e \right) \\
 &\quad + \varepsilon_{b[c]} \left(W_{nnd]a} - \frac{1}{2} k_{[d]a;n} + W_{d]nen} K_a{}^e \right) \\
 W_{0abc} &= 2W_{nnbc;\underline{a}} - 4W_{nnd[c} F_{b]a}{}^d + k_{a[b;c]} + f^d{}_{a[c} k_{b]d} \\
 W_{0a0b} &= \frac{-1}{n-2} W_{0abc;\underline{d}} \varepsilon^{cd} - \frac{1}{n-2} W_{0ecb} F^{ec}{}_a - \varepsilon^{cd} E_{0cd} \varepsilon_{ab} + E_{0ab} \\
 &\quad + \varepsilon^{cd} (E_{0acd;b} - E_{0acb;d}) + E_{0aec} f^{ec}{}_b - F_{abe} E^{ec}{}_c + F_{aec} E^{ec}{}_b \\
 &\quad + L^c W_{nbca} + K_{ab} E_{nc}{}^c - K_a{}^d E_{ndb}
 \end{aligned}
 \tag{A.25}$$

It follows from the relations above that the infinitesimal action of the structure group on W_{ijkl} is obtained from the appropriate chains of coframe derivatives of W_{anbn} .

REFERENCES

- [AG93] M. A. Akivis and V. V. Goldberg. *Projective differential geometry of submanifolds*, volume 49 of *North-Holland Mathematical Library*. North-Holland Publishing Co., Amsterdam, 1993. [11](#), [31](#)
- [AG96] M. A. Akivis and V. V. Goldberg. *Conformal differential geometry and its generalizations*. Pure and Applied Mathematics (New York). John Wiley & Sons Inc., New York, 1996. A Wiley-Interscience Publication. [2](#), [35](#), [38](#)
- [AJ16] A. B. Aazami and M. A. Javaloyes. Penrose’s singularity theorem in a Finsler spacetime. *Classical Quantum Gravity*, 33(2):025003, 22, 2016. [4](#)
- [Ale67] A. D. Alexandrov. A contribution to chronogeometry. *Canad. J. Math.*, 19:1119–1128, 1967. [2](#)
- [ANN15] I. Anderson, Z. Nie, and P. Nurowski. Non-rigid parabolic geometries of Monge type. *Adv. Math.*, 277:24–55, 2015. [4](#), [7](#), [21](#), [25](#)
- [AZ88] H. Akbar-Zadeh. Sur les espaces de Finsler à courbures sectionnelles constantes. *Acad. Roy. Belg. Bull. Cl. Sci. (5)*, 74(10):281–322, 1988. [6](#), [31](#)
- [AZ02] A. A. Agrachev and I. Zelenko. Geometry of Jacobi curves. I. *J. Dynam. Control Systems*, 8(1):93–140, 2002. [10](#), [13](#)
- [BCS00] D. Bao, S.-S. Chern, and Z. Shen. *An introduction to Riemann-Finsler geometry*, volume 200 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2000. [4](#), [5](#), [6](#), [31](#), [32](#)
- [BEE96] J. K. Beem, P. E. Ehrlich, and K. L. Easley. *Global Lorentzian geometry*, volume 202 of *Monographs and Textbooks in Pure and Applied Mathematics*. Marcel Dekker, Inc., New York, second edition, 1996. [27](#)
- [Bel01] F. A. Belgun. Null-geodesics in complex conformal manifolds and the LeBrun correspondence. *J. Reine Angew. Math.*, 536:43–63, 2001. [5](#)
- [Bes08] A. L. Besse. *Einstein manifolds*. Classics in Mathematics. Springer-Verlag, Berlin, 2008. Reprint of the 1987 edition. [38](#)
- [BRS04] D. Bao, C. Robles, and Z. Shen. Zermelo navigation on Riemannian manifolds. *J. Differential Geom.*, 66(3):377–435, 2004. [13](#)
- [Bry02] R. L. Bryant. Some remarks on Finsler manifolds with constant flag curvature. *Houston J. Math.*, 28(2):221–262, 2002. Special issue for S. S. Chern. [2](#), [10](#), [14](#), [27](#), [28](#)
- [Čap05] A. Čap. Correspondence spaces and twistor spaces for parabolic geometries. *J. Reine Angew. Math.*, 582:143–172, 2005. [5](#), [7](#)
- [Car30] É. Cartan. Sur un problème d’équivalence et la théorie des espaces métriques généralisés. *Mathematica*, 4:114–136, 1930. [4](#)
- [Car34] É. Cartan. *Les Espaces de Finsler, Exposés de Géométrie*. Paris, Hermann, 1934., 1934. [4](#), [28](#)
- [Che40] S.-S. Chern. The geometry of the differential equation $y''' = F(x, y, y', y'')$. *Sci. Rep. Nat. Tsing Hua Univ. (A)*, 4:97–111, 1940. [3](#)
- [Che48] S.-S. Chern. Local equivalence and Euclidean connections in Finsler spaces. *Sci. Rep. Nat. Tsing Hua Univ. Ser. A.*, 5:95–121, 1948. [4](#)
- [ČS00] A. Čap and H. Schichl. Parabolic geometries and canonical Cartan connections. *Hokkaido Math. J.*, 29(3):453–505, 2000. [22](#)
- [ČS09] A. Čap and J. Slovák. *Parabolic geometries. I*, volume 154 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2009. Background and general theory. [7](#), [20](#), [21](#)
- [DS01] A. Daigneault and A. Sangalli. Einstein’s static universe: an idea whose time has come back? *Notices Amer. Math. Soc.*, 48(1):9–16, 2001. [2](#)

- [DZ12] B. Doubrov and I. Zelenko. Prolongation of quasi-principal frame bundles and geometry of flag structures on manifolds. *arXiv preprint arXiv:1210.7334*, 2012. [5](#)
- [EPS72] J. Ehlers, F. A. E. Pirani, and A. Schild. The geometry of free fall and light propagation. *General relativity (papers in honour of J. L. Synge)*, pages 63–84, 1972. [2](#)
- [Fox05a] D. J. F. Fox. Contact path geometries. *arXiv preprint math/0508343*, 2005. [4](#), [5](#)
- [Fox05b] D. J. F. Fox. Contact projective structures. *Indiana Univ. Math. J.*, 54(6):1547–1598, 2005. [3](#)
- [Gar89] R. B. Gardner. *The method of equivalence and its applications*, volume 58 of *CBMS-NSF Regional Conference Series in Applied Mathematics*. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 1989. [7](#), [12](#), [41](#)
- [GH79] P. Griffiths and J. Harris. Algebraic geometry and local differential geometry. *Ann. Sci. École Norm. Sup. (4)*, 12(3):355–452, 1979. [15](#), [37](#)
- [GN09] M. Godlinski and P. Nurowski. Geometry of third-order odes. *arXiv preprint arXiv:0902.4129*, 2009. [3](#)
- [GPS05] A. García-Parrado and J. M. M. Senovilla. Causal structures and causal boundaries. *Classical Quantum Gravity*, 22(9):R1–R84, 2005. [2](#)
- [Gri83] P. A. Griffiths. *Exterior differential systems and the calculus of variations*, volume 25 of *Progress in Mathematics*. Birkhäuser, Boston, Mass., 1983. [5](#), [26](#)
- [Gro00] D. A. Grossman. Torsion-free path geometries and integrable second order ODE systems. *Selecta Math. (N.S.)*, 6(4):399–442, 2000. [4](#), [5](#), [6](#), [34](#), [39](#)
- [Gui89] V. Guillemin. *Cosmology in (2+1)-dimensions, cyclic models, and deformations of $M_{2,1}$* , volume 121 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 1989. [5](#)
- [Gut95] A. K. Guts. Axiomatic causal theory of space-time. *Gravitation and cosmology*, 1:301, 1995. [2](#)
- [HE73] S. W. Hawking and G. F. R. Ellis. *The large scale structure of space-time*. Cambridge University Press, London-New York, 1973. Cambridge Monographs on Mathematical Physics, No. 1. [2](#)
- [HÓ97] J. Hilgert and G. Ólafsson. *Causal symmetric spaces*, volume 18 of *Perspectives in Mathematics*. Academic Press, Inc., San Diego, CA, 1997. Geometry and harmonic analysis. [2](#)
- [HS10] J. Holland and G. Sparling. Causal geometries and third-order ordinary differential equations. *Arxiv preprint arXiv:1001.0202*, 2010. [3](#), [7](#)
- [HS11] J. Holland and G. Sparling. Causal geometries, null geodesics, and gravity. *arXiv preprint arXiv:1106.5254*, 2011. [3](#), [4](#), [27](#)
- [Hsu92] L. Hsu. Calculus of variations via the griffiths formalism. *Journal of Differential Geometry*, 36:551–589, 1992. [14](#), [26](#)
- [Hwa12] J.-M. Hwang. Geometry of varieties of minimal rational tangents. In *Current developments in algebraic geometry*, volume 59 of *Math. Sci. Res. Inst. Publ.*, pages 197–226. Cambridge Univ. Press, Cambridge, 2012. [5](#), [8](#)
- [Hwa13] J.-M. Hwang. Varieties of minimal rational tangents of codimension 1. *Ann. Sci. Éc. Norm. Supér. (4)*, 46(4):629–649 (2013), 2013. [5](#)
- [IL03] T. A. Ivey and J. M. Landsberg. *Cartan for beginners: differential geometry via moving frames and exterior differential systems*, volume 61 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2003. [12](#)
- [KP67] E. H. Kronheimer and R. Penrose. On the structure of causal spaces. *Proc. Cambridge Philos. Soc.*, 63:481–501, 1967. [2](#), [13](#)
- [KT14] B. Kruglikov and D. The. The gap phenomenon in parabolic geometries. *J. reine angew. Math.*, 2014. [21](#), [25](#)
- [LeB83] C. R. LeBrun. Spaces of complex null geodesics in complex-Riemannian geometry. *Trans. Amer. Math. Soc.*, 278(1):209–231, 1983. [5](#), [21](#)
- [LeB90] C. R. LeBrun. Twistors, ambitwistors, and conformal gravity. In *Twistors in mathematics and physics*, volume 156 of *London Math. Soc. Lecture Note Ser.*, pages 71–86. Cambridge Univ. Press, Cambridge, 1990. [5](#)
- [Lev93] A. V. Levichev. I. Segal’s chronometric theory as a completion of the special theory of relativity. *Izv. Vyssh. Uchebn. Zaved. Fiz.*, 36(8):84–89, 1993. [2](#)
- [LM87] P. Libermann and C.-M. Marle. *Symplectic Geometry and Analytical Mechanics*. D. Reidel Publishing Company, 1987. [12](#)
- [LM07] C. R. LeBrun and L. J. Mason. Nonlinear gravitons, null geodesics, and holomorphic disks. *Duke Math. J.*, 136(2):205–273, 2007. [38](#)
- [Low06] R. J. Low. The space of null geodesics (and a new causal boundary). In *Analytical and numerical approaches to mathematical relativity*, volume 692 of *Lecture Notes in Phys.*, pages 35–50. Springer, Berlin, 2006. [5](#)
- [Mak16] O. Makhmali. *Differential Geometric Aspects of Causal Structures*. PhD Thesis, McGill University, 2016. [7](#), [24](#), [25](#), [39](#)
- [Min15] E. Minguzzi. Raychaudhuri equation and singularity theorems in Finsler spacetimes. *Classical Quantum Gravity*, 32(18):185008, 26, 2015. [4](#)
- [Miy91] R. Miyaoka. Lie contact structures and normal Cartan connections. *Kodai Math. J.*, 14(1):13–41, 1991. [9](#)
- [Mor93] T. Morimoto. Geometric structures on filtered manifolds. *Hokkaido Math. J.*, 22(3):263–347, 1993. [22](#)
- [Olv95] P. J. Olver. *Equivalence, invariants, and symmetry*. Cambridge University Press, Cambridge, 1995. [7](#), [12](#), [41](#)

- [O’N83] B. O’Neill. *Semi-Riemannian geometry*, volume 103 of *Pure and Applied Mathematics*. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York, 1983. With applications to relativity. [26](#), [27](#)
- [Pen65] R. Penrose. Gravitational collapse and space-time singularities. *Phys. Rev. Lett.*, 14:57–59, 1965. [2](#)
- [Pen72] R. Penrose. *Techniques of differential topology in relativity*. Society for Industrial and Applied Mathematics, Philadelphia, Pa., 1972. Conference Board of the Mathematical Sciences Regional Conference Series in Applied Mathematics, No. 7. [2](#)
- [Pen76] R. Penrose. Nonlinear gravitons and curved twistor theory. *General Relativity and Gravitation*, 7(1):31–52, 1976. The riddle of gravitation—on the occasion of the 60th birthday of Peter G. Bergmann (Proc. Conf., Syracuse Univ., Syracuse, N. Y., 1975). [38](#)
- [Run83] H. Rund. Congruences of null-extremals in the calculus of variations. *J. Nat. Acad. Math. India*, 1(2):165–178, 1983. [5](#)
- [Run84] H. Rund. Null-distributions in the calculus of variations of multiple integrals. *Math. Colloq. Univ. Cape Town*, 13:57–81, 1984. [5](#)
- [Sac61] R. Sachs. Gravitational waves in general relativity. VI. The outgoing radiation condition. *Proc. Roy. Soc. Ser. A*, 264:309–338, 1961. [4](#)
- [Sas99] T. Sasaki. *Projective differential geometry and linear homogeneous differential equations*. Department of Mathematics, Kobe University, 1999. [11](#), [31](#), [37](#)
- [Sch13] P. Schapira. Hyperbolic systems and propagation on causal manifolds. *Lett. Math. Phys.*, 103(10):1149–1164, 2013. [2](#)
- [Seg76] I. E. Segal. *Mathematical cosmology and extragalactic astronomy*. Academic Press [Harcourt Brace Jovanovich, Publishers], New York-London, 1976. Pure and Applied Mathematics, Vol. 68. [2](#)
- [She01] Z. Shen. *Differential geometry of spray and Finsler spaces*. Kluwer Academic Publishers, Dordrecht, 2001. [5](#)
- [SY89] H. Sato and K. Yamaguchi. Lie contact manifolds. In *Geometry of manifolds (Matsumoto, 1988)*, volume 8 of *Perspect. Math.*, pages 191–238. Academic Press, Boston, MA, 1989. [9](#), [34](#)
- [SY98] H. Sato and A. Y. Yoshikawa. Third order ordinary differential equations and Legendre connections. *J. Math. Soc. Japan*, 50(4):993–1013, 1998. [3](#)
- [Tan79] N. Tanaka. On the equivalence problems associated with simple graded Lie algebras. *Hokkaido Math. J.*, 8(1):23–84, 1979. [3](#), [22](#)
- [Tau77] A. H. Taub. Book Review: Mathematical cosmology and extragalactic astronomy. *Bull. Amer. Math. Soc.*, 83(4):705–711, 1977. [2](#)
- [Wor84] L. I. Wormald. A critique of Segal’s chronometric theory. *Gen. Relativity Gravitation*, 16(4):393–401, 1984. [2](#)
- [Yam93] K. Yamaguchi. Differential systems associated with simple graded Lie algebras. In *Progress in differential geometry*, volume 22 of *Adv. Stud. Pure Math.*, pages 413–494. Math. Soc. Japan, Tokyo, 1993. [21](#), [22](#), [25](#)
- [Ye94] Y.-G. Ye. Extremal rays and null geodesics on a complex conformal manifold. *Internat. J. Math.*, 5(1):141–168, 1994. [5](#)
- [Zad09] V. Zadnik. Lie contact structures and chains. *arXiv preprint arXiv:0901.4433*, 2009. [35](#)
- [Zel09] I. Zelenko. On Tanaka’s prolongation procedure for filtered structures of constant type. *SIGMA Symmetry Integrability Geom. Methods Appl.*, 5:Paper 094, 21, 2009. [21](#)

DEPARTMENT OF MATHEMATICS AND STATISTICS, MCGILL UNIVERSITY, 805 SHERBROOKE STREET WEST, MONTREAL, QUEBEC, CANADA,, H3A 0B9

E-mail address: omid.makhmali@mail.mcgill.ca